

Multivariable (φ, Γ) -modules and smooth \mathfrak{o} -torsion representations

Gergely Zábrádi *

16th October 2018

Abstract

Let G be a \mathbb{Q}_p -split reductive group with connected centre and Borel subgroup $B = TN$. We construct a right exact functor D_Δ^\vee from the category of smooth modulo p^n representations of B to the category of projective limits of finitely generated étale (φ, Γ) -modules over a multivariable (indexed by the set of simple roots) commutative Laurent-series ring. These correspond to representations of a direct power of $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ via an equivalence of categories. Parabolic induction from a subgroup $P = L_P N_P$ gives rise to a basechange from a Laurent-series ring in those variables with corresponding simple roots contained in the Levi component L_P . D_Δ^\vee is exact and yields finitely generated objects on the category SP_A of finite length representations with subquotients of principal series as Jordan-Hölder factors. Lifting the functor D_Δ^\vee to all (noncommuting) variables indexed by the positive roots allows us to construct a G -equivariant sheaf $\mathfrak{Y}_{\pi, \Delta}$ on G/B and a G -equivariant continuous map from the Pontryagin dual π^\vee of a smooth representation π of G to the global sections $\mathfrak{Y}_{\pi, \Delta}(G/B)$. We deduce that D_Δ^\vee is fully faithful on the full subcategory of SP_A with Jordan-Hölder factors isomorphic to irreducible principal series.

Contents

1	Introduction	2
1.1	Background and motivation	2
1.2	Notations	4
1.3	Description of the results	5
2	Étale T_+-modules over $A((N_{\Delta, 0}))$	8
2.1	T_0 -invariant ideals in $A((N_{\Delta, 0}))$	8
2.2	A functor from smooth B -representations to étale T_+ -modules over $A((N_{\Delta, 0}))$.	10
2.3	The category $\mathcal{D}^{et}(T_+, A((N_{\Delta, 0})))$	11
2.4	A functor to usual (φ, Γ) -modules	12
2.5	Right exactness of D_Δ^\vee	14

*I would like to thank the MTA Alfréd Rényi Institute of Mathematics for its hospitality where parts of this work was written. This research was supported by a Hungarian OTKA Research grant K-100291 and by the János Bolyai Scholarship of the Hungarian Academy of Sciences.

3	Compatibility with tensor products and parabolic induction	16
3.1	Tensor products	16
3.2	Parabolic induction	20
3.3	Exactness of D_{Δ}^{\vee} on the category SP_A	25
4	Noncommutative theory	32
4.1	The ring $A((N_{\Delta, \infty}))$ and its first ring-theoretic properties	32
4.2	The equivalence of categories	35
4.3	A noncommutative variant of D_{Δ}^{\vee}	39
4.4	A natural transformation from the Schneider–Vigneras D -functor to $D_{\Delta, \infty}^{\vee}$	42
4.5	A G -equivariant sheaf on G/B attached to $D_{\Delta}^{\vee}(\pi)$	45
4.6	The fully faithful property of D_{Δ}^{\vee} for principal series	51

1 Introduction

1.1 Background and motivation

By now the p -adic Langlands correspondence for $\mathrm{GL}_2(\mathbb{Q}_p)$ is very well understood through the work of Berger [1], Breuil [4, 5, 6], Colmez [12], [13], Emerton [15], Kisin [21], Paškūnas [24] (see [7] for an overview). The starting point of Colmez’s work is Fontaine’s [18] theorem that the category of modulo p^h Galois representations of \mathbb{Q}_p is equivalent to the category of étale (φ, Γ) -modules over $\mathbb{Z}/p^h((X))$. One of Colmez’s breakthroughs was that he managed to relate smooth modulo p^h representations (therefore also continuous p -adic representations by letting $h \rightarrow \infty$ and inverting p) of $\mathrm{GL}_2(\mathbb{Q}_p)$ to (φ, Γ) -modules, too. The so-called “Montréal-functor” associates to a smooth mod p^h representation π of $\mathrm{GL}_2(\mathbb{Q}_p)$ (first restricting π to a Borel subgroup $B_2(\mathbb{Q}_p)$) an étale (φ, Γ) -module over $\mathbb{Z}/p^h((X))$. By Paškūnas’s work [24] this induces a bijection for certain p -adic Banach space representations of $\mathrm{GL}_2(\mathbb{Q}_p)$.

There have been attempts, for instance by Schneider and Vigneras [25], to generalize Colmez’s functor to other \mathbb{Q}_p -split reductive groups G . More recently, Breuil [8] (in a slightly more general setting allowing finite extensions of \mathbb{Q}_p , too) introduced a functor $D_{\xi}^{\vee} = D_{\xi, \ell}^{\vee}$ from smooth \mathbb{Z}/p^h -representations of G to projective limits of étale (φ, Γ) -modules. The construction depends on the choice of a cocharacter $\xi: \mathbf{G}_m \rightarrow T$ (with the property that the composition of ξ with all simple roots $\alpha \in \Delta$ is an isomorphism of \mathbf{G}_m) and on a Whittaker type functional ℓ from the unipotent radical N of a Borel subgroup $B = TN$ to \mathbb{Q}_p . In [8] (and also in [25]) ℓ is assumed to be *generic*, ie. ℓ induces an isomorphism $N_{\alpha} \rightarrow \mathbb{Q}_p$ for the root subgroups N_{α} of all simple roots $\alpha \in \Delta$ with respect to B . The action of φ (resp. of $\Gamma \cong \mathbb{Z}_p^{\times}$) on $D_{\xi}^{\vee}(\pi)$ for a smooth mod p^h representation π of G comes from the (inverse of the) action of $\xi(p)$ (resp. of $\xi(\mathbb{Z}_p^{\times})$) on π . The functor $D_{\xi, \ell}^{\vee}$ has very promising properties: it is right exact and compatible with tensor products and with parabolic induction. Moreover, $D_{\xi, \ell}^{\vee}$ is exact and produces finitely generated objects on the category SP_A of finite length representations with all Jordan-Hölder factors appearing as a subquotient of principal series representations (ie. of $\mathrm{Ind}_B^G \chi$ for some character χ of T). Finally, $D_{\xi, \ell}^{\vee}$ is compatible with the conjectures in [10] made from a global point of view. The assumption on the genericity of ℓ is needed crucially for some of these properties, in particular for the exactness on SP_A and for the compatibility with [10]. However, if ℓ is a generic Whittaker functional then the

functor $D_{\xi,\ell}^\vee$ loses a lot of information, one cannot possibly recover the representation π from the attached (φ, Γ) -module $D_{\xi,\ell}^\vee(\pi)$ (by the methods developed in [26] or otherwise). This has also been predicted by the work of Breuil and Paškūnas [11]: when one moves beyond $\mathrm{GL}_2(\mathbb{Q}_p)$ then there are much more representations on the automorphic side than on the Galois side. So if we would like to have a bijection for some large class of representations on the reductive group side, we need to put additional data on our Galois-representations. One candidate is that we could perhaps equip the Galois representation with an additional character of the torus $T/\xi(\mathbb{Q}_p^\times)$ extending the action of φ and Γ . The heuristics for this is that even in the case of $\mathrm{GL}_2(\mathbb{Q}_p)$ a central character appears naturally on the attached (φ, Γ) -module. However, if ℓ is generic then the action of φ and Γ on $D_{\xi,\ell}^\vee(\pi)$ cannot be extended to the dominant submonoid $T_+ \subset T$ since in this case the kernel $H_{gen} = \mathrm{Ker}(\ell: N \rightarrow \mathbb{Q}_p)$ is not invariant under the conjugation action of any larger subgroup of T than the product of the image of ξ and the centre. On the other hand, if we choose ℓ to be very far from being generic, ie. $\ell = \ell_\alpha$ is the projection onto a root subgroup N_α for some simple root $\alpha \in \Delta$ then we do have an additional action of T_+ on $D_{\xi,\ell}^\vee(\pi)$ as shown by the present author and Erdélyi [17]. Moreover, in op. cit. a natural transformation $\beta_{G/B,\cdot}$ from the functor $\pi \mapsto \pi^\vee$ (taking Pontryagin duals) to the global sections $\mathfrak{Y}_{\alpha,\pi}(G/B)$ of a G -equivariant sheaf $\mathfrak{Y}_{\alpha,\pi}$ on the flag variety G/B associated to the étale T_+ -module $D_{\xi,\ell}^\vee(\pi)$ is constructed for the choice $\ell = \ell_\alpha$. The map $\beta_{G/B,\pi}: \pi^\vee \rightarrow \mathfrak{Y}_{\alpha,\pi}(G/B)$ is nonzero whenever $D_{\xi,\ell}^\vee(\pi)$ is nonzero. However, as mentioned above, for non-generic ℓ the functor $D_{\xi,\ell}^\vee$ does not have so good exactness and compatibility properties.

The goal of this paper is to combine all the mentioned good properties of the above approaches. In order to do this we are going to use *multivariable* (φ, Γ) -modules in the variables X_α ($\alpha \in \Delta$). More concretely, consider the Laurent series ring $\mathbb{Z}/p^h((X_\alpha \mid \alpha \in \Delta)) := \mathbb{Z}/p^h[[X_\alpha, \alpha \in \Delta]][X_\alpha^{-1} \mid \alpha \in \Delta]$ with the conjugation action of the monoid $T_+ := \{t \in T \mid \alpha(t) \in \mathbb{Z}_p \text{ for all } \alpha \in \Delta\}$. In an analogous way to [8] we construct a functor D_Δ^\vee from smooth mod p^h -representations of a \mathbb{Q}_p -split connected reductive group G with connected centre to the category of projective limits of finitely generated étale T_+ -modules over $\mathbb{Z}/p^h((X_\alpha \mid \alpha \in \Delta))$. Moreover, in [29] a pair \mathbb{D} and \mathbb{V} of quasi-inverse equivalences of categories is constructed between the category of continuous mod p^n representations of the $|\Delta|$ th direct power of the Galois group $\mathrm{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ (endowed with a character of \mathbb{Q}_p^\times) and multivariable étale T_+ -modules. One can pass to usual (φ, Γ) -modules by identifying the variables X_α with each other—this step corresponds to the restriction of a representation of $\mathrm{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)^{|\Delta|}$ to the diagonal embedding of $\mathrm{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ (Cor. 3.10 in [29]). When doing so we must forget the action of the monoid T_+ just keeping the action of $\varphi^{\mathbb{N}}\Gamma = \xi(\mathbb{Z}_p \setminus \{0\}) \subset T_+$ (or possibly also the action of the centre of G) as the kernel $(X_\alpha - X_\beta \mid \alpha, \beta \in \Delta)$ of this identification is not stable under T_+ . This assignment is faithful and exact in general, but definitely not full. In all known cases—including objects in the category SP_A and parabolically induced representations from the product of copies of $\mathrm{GL}_2(\mathbb{Q}_p)$ and a torus—the resulting Galois representation will coincide with Breuil’s $\mathbb{V}_F \circ D_{\xi,\ell}^\vee(\pi)$ (for generic ℓ) where \mathbb{V}_F stands for Fontaine’s equivalence. Whether or not this is true in general is an open question. If $P = L_P N_P$ is a standard parabolic subgroup with Levi component L_P isomorphic to the product of copies of $\mathrm{GL}_2(\mathbb{Q}_p)$ and a torus, then the value of $\mathbb{V} \circ D_\Delta^\vee$ at parabolically induced representations $\mathrm{Ind}_P^G \pi_P$ is well-described in terms of tensor product of Galois representations for each $\alpha \in \Delta$. It can be shown using the fully faithful property of \mathbb{D} that the resulting

multivariable (φ, Γ) -modules $D_\Delta^\vee(\text{Ind}_P^G \pi_P)$ are therefore pairwise non-isomorphic for all the irreducible mod p representations of G arising this way (for varying P of this form).

Apart from all the above mentioned exactness and compatibility properties D_Δ^\vee has the following additional features: induction from a parabolic subgroup $P = L_P N_P$ corresponds to basechange from $\mathbb{Z}/(p^n)((X_\alpha \mid \alpha \in \Delta_P))$ to $\mathbb{Z}/p^h((X_\alpha \mid \alpha \in \Delta))$ where $\Delta_P \subseteq \Delta$ consists of those simple roots whose root subgroups are contained in the Levi component L_P . On the Galois side this means, in particular, that the copy of the Galois group $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ corresponding to those simple roots $\alpha \in \Delta$ whose root subgroup is not contained in the Levi L_P acts on $\mathbb{V} \circ D_\Delta^\vee(\text{Ind}_P^G \pi_P)$ via a character. This could hopefully lead to detecting P from the attached T_+ -module over $\mathbb{Z}/p^h((X_\alpha \mid \alpha \in \Delta))$. Another promising property of D_Δ^\vee is that we can indeed recover successive extensions π of irreducible principal series representations from $D_\Delta^\vee(\pi)$. In other words we show—using the methods of [26] [17] realizing π^\vee as a G -invariant subspace of the global sections of a G -equivariant sheaf on G/B —that D_Δ^\vee is fully faithful on the category SP_A^0 of these representations. By the aforementioned work of Breuil and Paškūnas [11] we cannot expect a bijection between smooth \mathbb{Z}/p^h -representations of G and mod p^h Galois representations of \mathbb{Q}_p . However, this work could be considered as evidence that there might still be a bijection between a large class of smooth \mathbb{Z}/p^h -representations of G and certain representations of $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)^{|\Delta|}$.

Moreover, Breuil, Herzig, and Schraen [9] predict that Breuil’s functor $\mathbb{V}_F \circ D_{\xi, \ell}^\vee(\pi)$ on a representation of $\text{GL}_n(\mathbb{Q}_p)$ built out from some mod p Hecke isotopic subspace would give something like an “internal” tensor product $\rho \otimes_{\mathbb{F}_p} \wedge^2(\rho) \otimes_{\mathbb{F}_p} \cdots \otimes_{\mathbb{F}_p} \wedge^{n-1}(\rho)$ for some local Galois representation ρ of dimension n furnished by the global theory. Now the functor $\mathbb{V} \circ D_\Delta^\vee$ should give the same, but “external” tensor product, instead of internal (ie. different copies of $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ for each term in the tensor product). This could perhaps explain why the individual $\wedge^i(\rho)$ appear in the Shimura cohomology of unitary groups of type $U(i, n-i)$, but not their internal tensor product.

Another motivation is that the Robba versions of multivariable (φ, Γ) -modules seem to play a role [2] [3] [20] in the case of the p -adic Langlands programme for $\text{GL}_2(F)$ for finite extensions $F \neq \mathbb{Q}_p$, too.

1.2 Notations

Let $G = \mathbf{G}(\mathbb{Q}_p)$ be the \mathbb{Q}_p -points of a \mathbb{Q}_p -split connected reductive group \mathbf{G} defined over \mathbb{Z}_p with connected centre and a fixed split Borel subgroup $\mathbf{B} = \mathbf{TN}$. Put $B := \mathbf{B}(\mathbb{Q}_p)$, $T := \mathbf{T}(\mathbb{Q}_p)$, and $N := \mathbf{N}(\mathbb{Q}_p)$. We denote by Φ^+ the set of roots of T in N , by $\Delta \subset \Phi^+$ the set of simple roots, and by $u_\alpha : \mathbb{G}_a \rightarrow N_\alpha$, for $\alpha \in \Phi^+$, a \mathbb{Q}_p -homomorphism onto the root subgroup N_α of N such that $tu_\alpha(x)t^{-1} = u_\alpha(\alpha(t)x)$ for $x \in \mathbb{Q}_p$ and $t \in T(\mathbb{Q}_p)$, and $N_0 = \prod_{\alpha \in \Phi^+} u_\alpha(\mathbb{Z}_p)$ is a compact open subgroup of $N(\mathbb{Q}_p)$. We put $n_\alpha := u_\alpha(1)$ and $N_{\alpha,0} := u_\alpha(\mathbb{Z}_p)$ for the image of u_α on \mathbb{Z}_p . We denote by T_+ the monoid of dominant elements t in T such that $\text{val}_p(\alpha(t)) \geq 0$ for all $\alpha \in \Phi^+$, by $T_0 \subset T_+$ the maximal subgroup, and we put $B_+ = N_0 T_+$, $B_0 = N_0 T_0$.

Let K be a finite extension of \mathbb{Q}_p with ring of integers \mathfrak{o} , uniformizer ϖ , and residue field $\kappa := \mathfrak{o}/\varpi$. By a smooth \mathfrak{o} -torsion representation π of G (resp. of B) we mean a torsion \mathfrak{o} -module π together with a smooth (ie. stabilizers are open) and linear action of the group G (resp. of B). We will consider representations π with $\varpi^h \pi = 0$ for some $h \geq 1$ and put

$A := o/\varpi^h$.

The natural conjugation action of T_+ on N_0 extends to an action on the Iwasawa A -algebra $A[[N_0]]$. For $t \in T_+$ we denote this action of t on $A[[N_0]]$ by φ_t . The map $\varphi_t: A[[N_0]] \rightarrow A[[N_0]]$ is an injective ring homomorphism with a distinguished left inverse $\psi_t: A[[N_0]] \rightarrow A[[N_0]]$ satisfying $\psi_t \circ \varphi_t = \text{id}_{A[[N_0]]}$ and $\psi_t(u\varphi_t(\lambda)) = \psi_t(\varphi_t(\lambda)u) = 0$ for all $u \in N_0 \setminus tN_0t^{-1}$ and $\lambda \in A[[N_0]]$. Further, the normal subgroup $H_{\Delta,0} := \prod_{\beta \in \Phi^+ \setminus \Delta} N_{\beta,0}$ is invariant under the action of T_+ so the quotient group $N_{\Delta,0} := N_0/H_{\Delta,0} \cong \prod_{\alpha \in \Delta} N_{\alpha,0}$ also inherits the action of T_+ . The Iwasawa algebra $A[[N_{\Delta,0}]]$ can be identified with the multivariable power series ring $A[[X_\alpha \mid \alpha \in \Delta]]$ by the map $n_\alpha - 1 \mapsto X_\alpha$ ($\alpha \in \Delta$). We define $A((N_{\Delta,0}))$ as the localization $A[[N_{\Delta,0}]] [X_\alpha^{-1}, \alpha \in \Delta]$. We also denote by $\varphi_t: A[[N_{\Delta,0}]] \rightarrow A[[N_{\Delta,0}]]$ (resp. $\varphi_t: A((N_{\Delta,0})) \rightarrow A((N_{\Delta,0}))$) the induced action of $t \in T_+$ on these rings. By an étale T_+ -module over $A((N_{\Delta,0}))$ we mean a (unless otherwise mentioned) finitely generated module M over $A((N_{\Delta,0}))$ together with a semilinear action of the monoid T_+ (also denoted by φ_t for $t \in T_+$) such that the maps

$$\text{id} \otimes \varphi_t: \varphi_t^* M := A((N_{\Delta,0})) \otimes_{A((N_{\Delta,0}), \varphi_t)} M \rightarrow M$$

are isomorphisms for all $t \in T_+$.

Since the centre of G is assumed to be connected, there exists a cocharacter $\lambda_{\alpha^\vee}: \mathbb{Q}_p^\times \rightarrow T$ such that $\alpha \circ \lambda_{\alpha^\vee}$ is the identity on \mathbb{Q}_p^\times for each $\alpha \in \Delta$ and $\beta \circ \lambda_{\alpha^\vee} = 1$ for all $\beta \neq \alpha \in \Delta$. Note that λ_{α^\vee} is only unique up to a cocharacter of the centre $Z(G)$. We put $\xi := \sum_{\alpha \in \Delta} \lambda_{\alpha^\vee}$, $\Gamma := \xi(\mathbb{Z}_p^\times) \leq T$, and often denote the action of $s := \xi(p)$ by $\varphi = \varphi_s$. Further, for each $\alpha \in \Delta$ we set $t_\alpha := \lambda_{\alpha^\vee}(p)$.

For example, $\mathbf{G} = \text{GL}_n$, B is the subgroup of upper triangular matrices, N consists of the strictly upper triangular matrices (1 on the diagonal), T is the diagonal subgroup, $N_0 = \mathbf{N}(\mathbb{Z}_p)$, the simple roots are $\alpha_1, \dots, \alpha_{n-1}$ where $\alpha_i(\text{diag}(t_1, \dots, t_n)) = t_i t_{i+1}^{-1}$, $u_{\alpha_i}(\cdot)$ is the strictly upper triangular matrix, with $(i, i+1)$ -coefficient \cdot and 0 everywhere else.

For a finite index subgroup \mathcal{G}_2 in a group \mathcal{G}_1 we denote by $J(\mathcal{G}_1/\mathcal{G}_2) \subset \mathcal{G}_1$ a (fixed) set of representatives of the left cosets in $\mathcal{G}_1/\mathcal{G}_2$.

1.3 Description of the results

In section 2 we describe the first properties of étale T_+ -modules over $A((N_{\Delta,0}))$ (the “multivariable (φ, Γ) -modules” in the title). Even though the ring $A((N_{\Delta,0}))$ is not artinian, the existence of an action of T_0 improves its properties: by the nonexistence of T_0 -invariant ideals in $\kappa((N_{\Delta,0}))$ it follows that any finitely generated module over $A((N_{\Delta,0}))$ admitting a semilinear action of T_0 has finite length (in the category of modules with semilinear T_0 -action). This fact allows us to construct a functor D_Δ^\vee from the category of smooth A -representations of the Borel B to projective limits of finitely generated étale T_+ -modules over $A((N_{\Delta,0}))$ in an analogous way to Breuil’s functor [8]. More precisely, we consider the skew polynomial ring $A[[N_{\Delta,0}]] [F_\alpha \mid \alpha \in \Delta]$ where the variables F_α commute with each other and we have $F_\alpha \lambda = (t_\alpha \lambda t_\alpha^{-1}) F_\alpha$ for $\lambda \in A[[N_{\Delta,0}]]$. For a smooth representation π of B over A we denote by $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ the set of finitely generated $A[[N_{\Delta,0}]] [F_\alpha \mid \alpha \in \Delta]$ -submodules of $\pi^{H_{\Delta,0}}$ that are stable under the action of T_0 and admissible as a representation of $N_{\Delta,0} = N_0/H_{\Delta,0}$. Here F_α acts on $\pi^{H_{\Delta,0}}$ by the Hecke action of $t_\alpha \in T_+$, ie. $F_\alpha v := \text{Tr}_{H_{\Delta,0}/t_\alpha H_{\Delta,0} t_\alpha^{-1}}(t_\alpha v)$ for $v \in \pi^{H_{\Delta,0}}$.

Then the functor D_Δ^\vee is defined by the projective limit

$$D_\Delta^\vee(\pi) := \varprojlim_{M \in \mathcal{M}_\Delta(\pi^{H_\Delta, 0})} M^\vee[1/X_\Delta]$$

where $X_\Delta = \prod_{\alpha \in \Delta} X_\alpha$ is the product of all the variables $X_\alpha = n_\alpha - 1$ in the power series ring $A[[N_{\Delta, 0}]]$.

If we define

$$\ell: N \rightarrow N/[N, N] = \prod_{\alpha \in \Delta} N_\alpha \xrightarrow{\sum_{\alpha \in \Delta} u_\alpha^{-1}} \mathbb{Q}_p$$

in a generic way and extend this to the Iwasawa algebra $A[[N_{\Delta, 0}]]$ then we find that $\ell(X_\alpha) = X$ for all $\alpha \in \Delta$ after the identification $A[[\mathbb{Z}_p]] \cong A[[X]]$. Therefore we may extend ℓ to a map $\ell: A((N_{\Delta, 0})) \rightarrow A((X))$ of Laurent series rings. Note that the kernel of ℓ is not stable under the action of T_+ , but it is stable under the action of φ and Γ . So we obtain a reduction map $A((X)) \otimes_{A((N_{\Delta, 0})), \ell} \cdot$ from étale T_+ -modules to usual étale (φ, Γ) -modules. We show that this reduction map is faithful and exact which implies

Theorem A. *The functor D_Δ^\vee is right exact.*

In particular, one has a natural transformation from Breuil's functor $D_{\xi, \ell}^\vee$ to the composite $A((X)) \otimes_{A((N_{\Delta, 0})), \ell} D_\Delta^\vee$. When restricted to the category SP_A , this is an isomorphism. Moreover, this is also an isomorphism for objects obtained by parabolic induction from a subgroup with Levi component isomorphic to the product of copies of $\mathrm{GL}_2(\mathbb{Q}_p)$ and a split torus.

Section 3 is devoted to various compatibility results. The first is the compatibility with products $G \times G'$ of groups with simple roots Δ , resp. Δ' . The value of $D_{\Delta \cup \Delta'}^\vee$ on a tensor product $\pi \otimes_\kappa \pi'$ of representations π of G (resp. π' of G') is the completed tensor product $D_\Delta^\vee(\pi) \hat{\otimes}_\kappa D_{\Delta'}^\vee(\pi')$. Note that this is a module over a multivariate Laurent series ring $A((N_{\Delta \cup \Delta', 0}))$ in variables indexed by the union $\Delta \cup \Delta'$. Similarly, we have a compatibility result for parabolic induction: Let $P = L_P N_P$ be a parabolic subgroup containing B and π_P a smooth representation of L_P over A viewed as representation of the opposite parabolic P^- . Denote by $\Delta_P \subseteq \Delta$ the set of those simple roots whose root subgroups are contained in the Levi component L_P . We show

Theorem B. *Let π_P be a smooth locally admissible representation of L_P over A which we view by inflation as a representation of P^- . We have an isomorphism*

$$D_\Delta^\vee(\mathrm{Ind}_{P^-}^G \pi_P) \cong A((N_{\Delta, 0})) \hat{\otimes}_{A((N_{\Delta_P, 0}))} D_{\Delta_P}^\vee(\pi_P)$$

in the category $\mathcal{D}^{\mathrm{pro-et}}(T_+, A((N_{\Delta, 0})))$.

On one hand, the above result shows that D_Δ^\vee is nonzero and finitely generated on parabolically induced representations from products of copies of $\mathrm{GL}_2(\mathbb{Q}_p)$ and a torus unless one of the representations of $\mathrm{GL}_2(\mathbb{Q}_p)$ is finite dimensional. Moreover, combined with the right exactness we also know this for extensions of representations of this type just like for Breuil's functor [8]. On the other hand, this might lead to another characterization of supercuspidal representations: it would be natural to expect that if π is an irreducible supercuspidal representation then $D_\Delta^\vee(\pi)$ cannot be induced from a T_+ -module in less variables. However, showing this would require a better understanding of supercuspidals beyond GL_2 .

Let SP_A be the category of smooth finite length representations of G whose Jordan-Hölder factors are subquotients of principal series. We end section 3 by showing

Theorem C. *The restriction of D_Δ^\vee to SP_A is exact and produces finitely generated objects.*

The proof of this builds on showing that the finitely generated $A[[N_{\Delta,0}]] [F_\alpha \mid \alpha \in \Delta]$ -submodules of representations in SP_A are in fact finitely presented. This substitutes the arguments using the coherence [14] of the one-variable analogue $A[[X]][F]$ in the classical GL_2 -situation as the ring $A[[N_{\Delta,0}]] [F_\alpha \mid \alpha \in \Delta]$ is apparently not coherent.

In section 4 we develop a noncommutative analogue of D_Δ^\vee as in [17] for Breuil's functor. The first step is the construction of the ring $A((N_{\Delta,\infty}))$ as a projective limit $\varprojlim_k A((N_{\Delta,k}))$ where the finite layers $A((N_{\Delta,k})) := A[[N_{\Delta,k}]] [\varphi_s^{kn_0}(X_\alpha)^{-1}]$ (where $n_0 = n_0(G) \in \mathbb{N}$ is the maximum of the degrees of the algebraic characters $\beta \circ \xi: \mathbf{G}_m \rightarrow \mathbf{G}_m$ for all positive roots $\beta \in \Phi^+$) are defined as localisations of the Iwasawa algebra $A[[N_{\Delta,k}]]$. Here the group $N_{\Delta,k} := N_0/H_{\Delta,k}$ is the extension of $N_{\Delta,0}$ by a finite p -group $H_{\Delta,0}/H_{\Delta,k}$ where $H_{\Delta,k}$ is the smallest normal subgroup in N_0 containing $s^k H_{\Delta,0} s^{-k}$. Note that unlike in the one variable localization $\Lambda_\ell(N_0)$ we do not have a section of the group homomorphism $N_{\Delta,k} \rightarrow N_{\Delta,0}$. However, restricting to the image of the conjugation by s^{kn_0} , we do: this allows us to build a functor $\mathbb{M}_{k,0}$ from the category $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ of finitely generated étale T_+ -modules over $A((N_{\Delta,0}))$ to the category $\mathcal{D}^{et}(T_+, A((N_{\Delta,k})))$ of finitely generated étale T_+ -modules over $A((N_{\Delta,k}))$. Putting $\mathbb{M}_{\infty,0} := \varprojlim_k \mathbb{M}_{k,0}$ and $\mathbb{D}_{0,\infty}$ to be the functor from the category $\mathcal{D}^{et}(T_+, A((N_{\Delta,\infty})))$ of finitely generated étale T_+ -modules over $A((N_{\Delta,\infty}))$ to $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ induced by the reduction map $A((N_{\Delta,\infty})) \rightarrow A((N_{\Delta,0}))$ we obtain

Theorem D. *The functors $\mathbb{M}_{\infty,0}$ and $\mathbb{D}_{0,\infty}$ are quasi-inverse equivalences of categories between $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ and $\mathcal{D}^{et}(T_+, A((N_{\Delta,\infty})))$.*

By considering finitely generated $A[[N_{\Delta,k}]] [F_{\alpha,k} \mid \alpha \in \Delta]$ -submodules of $\pi^{H_{\Delta,k}}$ that are stable under the action of T_0 and are admissible as representations of $N_{\Delta,k}$ we introduce the functors $D_{\Delta,k}^\vee$ analogous to D_Δ^\vee for all $k \geq 0$ and we put $D_{\Delta,\infty}^\vee(\pi) := \varprojlim_k D_{\Delta,k}^\vee(\pi)$ for a smooth representation π of B over A . This corresponds to $D_\Delta^\vee(\pi)$ via the extension of the equivalence of categories in Theorem D to pro-objects on both sides. The universal property of $D_{\Delta,\infty}^\vee$ leads to its alternative description via the Schneider–Vigneras functor $D_{SV}(\pi)$ (and via its étale hull $\widetilde{D}_{SV}(\pi)$):

Theorem E. *We have*

$$D_{\Delta,\infty}^\vee(\pi) \cong \varprojlim_D D$$

where D runs through the finitely generated étale T_+ -modules over $A((N_{\Delta,\infty}))$ arising as a quotient of $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D}_{SV}(\pi)$ such that the quotient map is continuous in the weak topology of D and the final topology on $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D}_{SV}(\pi)$ of the map $1 \otimes \iota: D_{SV}(\pi) \rightarrow A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D}_{SV}(\pi)$.

Finally, we turn to the question of reconstructing the smooth representation π of G from $D_\Delta^\vee(\pi)$. This is certainly not possible in general, as for instance finite dimensional representations are in the kernel of D_Δ^\vee (unless the set Δ of simple roots is empty). However, using the ideas of [26] we show the following positive results in this direction. For an object $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ we denote by $\widetilde{M}_\infty^\vee$ the étale hull of the image M_∞^\vee of the natural map from π^\vee to the étale T_+ -module $\mathbb{M}_{\infty,0}(M^\vee[1/X_\Delta])$.

Theorem F. *For any smooth o -torsion representation π of G and any $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ there exists a G -equivariant sheaf $\mathfrak{Y}_{\pi,M}$ on G/B with sections $\mathfrak{Y}_{\pi,M}(\mathcal{C}_0)$ on \mathcal{C}_0 isomorphic to $\widetilde{M}_\infty^\vee$ as an étale T_+ -module over $A[[N_0]]$. Moreover, we have a G -equivariant continuous map $\beta_{G/B,M}$ from the Pontryagin dual π^\vee to the global sections $\mathfrak{Y}_{\pi,M}(G/B)$ that is natural in both π and M , and is nonzero unless $M^\vee[1/X_\Delta] = 0$.*

Here we in fact use the G -action on π in order to construct the sheaf $\mathfrak{Y}_{\pi,M}$ unlike in [26] where the operators $\mathcal{H}_g = \text{res}(g\mathcal{C}_0 \cap \mathcal{C}_0) \circ (g \cdot)$ for the open cell $\mathcal{C}_0 := N_0\overline{B} \subset G/\overline{B} \cong G/B$ are constructed as a limit. Apparently the formulas defining this limit do not converge in the weak topology of the finitely generated $A((N_{\Delta,\infty}))$ -module $M_\infty^\vee[1/X_\Delta]$. Nevertheless, if π is irreducible and $D_\Delta^\vee(\pi) \neq 0$ then we can realize π^\vee as a subrepresentation of the global sections of a G -equivariant sheaf on G/B whose space of sections on \mathcal{C}_0 is “small” in the sense that it is contained in a finitely generated $A((N_{\Delta,\infty}))$ -module. Let us denote by SP_A^0 the full subcategory of SP_A containing those representations whose Jordan-Hölder factors are irreducible principle series. As an application of the methods above we prove

Theorem G. *The restriction of D_Δ^\vee to the category SP_A^0 is fully faithful.*

In particular, the forgetful functor restricting π to B is also fully faithful on SP_A^0 as D_Δ^\vee factors through this.

Acknowledgements

My debt to the works of Christophe Breuil [8], Pierre Colmez [12] [13], Peter Schneider, and Marie-France Vignéras [25] [26] will be obvious to the reader. I would also like to thank Márton Erdélyi, Jan Kohlhaase, Vytautas Paškūnas, Peter Schneider, and Tamás Szamuely for discussions on the topic. I am grateful to the referee for the careful reading of the manuscript and for their various comments.

2 Étale T_+ -modules over $A((N_{\Delta,0}))$

Since the centre of G is assumed to be connected, there exists a system $(\lambda_{\alpha^\vee})_{\alpha \in \Delta} \in X^\vee(T)^\Delta$ of cocharacters with the property $\beta \circ \lambda_{\alpha^\vee} = 1$ for all $\alpha \neq \beta \in \Delta$ and $\alpha \circ \lambda_{\alpha^\vee} = \text{id}_{\mathbf{G}_m}$. As in [8] and [25] we put $\xi := \sum_{\alpha \in \Delta} \lambda_{\alpha^\vee}$. Further, we put $t_\alpha := \lambda_{\alpha^\vee}(p) \in T$ for each $\alpha \in \Delta$ and denote by φ_α the conjugation action of t_α on $A[[N_{\Delta,0}]]$ and on $A((N_{\Delta,0}))$. By definition we have $s = \xi(p) = \prod_{\alpha \in \Delta} t_\alpha$. We form the skew-polynomial ring $A[[N_{\Delta,0}]] [F_\Delta] := A[[N_{\Delta,0}]] [F_\alpha \mid \alpha \in \Delta]$ in the variables F_α ($\alpha \in \Delta$) that commute with each other and satisfy $F_\alpha \lambda = \varphi_\alpha(\lambda) F_\alpha$ for any $\lambda \in \kappa[[N_{\Delta,0}]]$. Note that we may extend the conjugation action of the group T_0 on $A[[N_{\Delta,0}]]$ to the ring $A[[N_{\Delta,0}]] [F_\Delta]$ by acting trivially on the variables F_α ($\alpha \in \Delta$). Note that T_0 and the elements t_α ($\alpha \in \Delta$) generate T_+ under our assumption that the centre of G is connected.

2.1 T_0 -invariant ideals in $A((N_{\Delta,0}))$

Proposition 2.1. *The ring $\kappa((N_{\Delta,0}))$ does not have any nontrivial T_0 -invariant ideals.*

Proof. By our assumption that G has connected centre the group homomorphisms $\mathbb{Z}_p^\times \xrightarrow{\lambda_{\alpha^\vee}} T_0$ give rise to a subgroup $T_{\Delta,0} := \prod_{\alpha \in \Delta} \lambda_{\alpha^\vee}(\mathbb{Z}_p^\times) \leq T_0$. Note that this product is direct since $T_{\alpha,0} := \lambda_{\alpha^\vee}(\mathbb{Z}_p^\times)$ is contained in the kernel of β for each $\alpha \neq \beta \in \Delta$. So we have an action of $(\mathbb{Z}_p^\times)^\Delta$ on $\kappa((N_{\Delta,0})) = \kappa((X_\alpha \mid \alpha \in \Delta))$ such that the copy of \mathbb{Z}_p^\times indexed by α acts naturally on the corresponding variable X_α by the formula $\gamma \in \mathbb{Z}_p^\times: X_\alpha \mapsto (1 + X_\alpha)^\gamma - 1$ and trivially on all the other variables X_β for all pairs $\beta \neq \alpha \in \Delta$. We prove the (formally stronger) statement that $\kappa((N_{\Delta,0}))$ does not have any $T_{\Delta,0}$ -invariant ideals by induction on the cardinality of the set Δ . For $|\Delta| = 1$ the statement is trivial since $\kappa((X))$ is a field. Now assume the statement for $|\Delta| < r$ for some $1 < r$ choose a $T_{\Delta,0}$ -invariant ideal $0 \neq I \triangleleft \kappa((N_{\Delta,0}))$ with $|\Delta| = r$. Put $J := I \cap \kappa[[N_{\Delta,0}]] \triangleleft \kappa[[N_{\Delta,0}]]$ and choose any $\alpha \in \Delta$. Any element $\mu \in \kappa[[N_{\Delta,0}]]$ can uniquely be written as an infinite sum $\mu = \sum_{j=0}^{\infty} \mu_j X_\alpha^j$ with $\mu_j \in \kappa[[N_{\Delta \setminus \{\alpha\},0}]]$ ($j \geq 0$) where $N_{\Delta \setminus \{\alpha\},0} = \prod_{\beta \neq \alpha \in \Delta} N_{\beta,0}$. We define

$$J_{\alpha,i} := \{ \lambda \in \kappa[[N_{\Delta \setminus \{\alpha\},0}]] \mid \exists \mu = \sum_{j=0}^{\infty} \mu_j X_\alpha^j \in J, \text{ s. t. } \mu_i = \lambda \}$$

for each $i \geq 0$. These are ideals in $\kappa[[N_{\Delta \setminus \{\alpha\},0}]]$.

Lemma 2.2. *We have $J_{\alpha,0} = J_{\alpha,1} = \dots = J_{\alpha,i} = \dots$.*

Proof. The inclusions $J_{\alpha,0} \subseteq J_{\alpha,1} \subseteq \dots \subseteq J_{\alpha,i} \subseteq \dots$ are clear (we can multiply an element in J by X_α). Conversely, assume that $J_{\alpha,0} \subsetneq J_{\alpha,i}$ for some integer $i > 0$. Assume $i > 0$ is minimal with this property and choose an element $\mu \in J$ with $\mu_i \in J_{\alpha,i} \setminus J_{\alpha,0}$. Assume there is an index $j > 0$ such that $\mu_j \neq 0$ and $\mu_j \in J_{\alpha,0}$, and choose a $\nu_j \in J$ with $\nu_{j,0} = \mu_j$. Then $\mu' := \mu - X_\alpha^j \nu_j$ also lies in J and has the property that $\mu'_i \notin J_{\alpha,0}$. Indeed, if $i < j$ then this is clear. Otherwise by the minimality of i , the coefficient of X_α^{i-j} in ν_j lies in $J_{\alpha,0}$ (and is equal to $\mu_i - \mu'_i$ for $i \geq j$). Since any $\kappa[[N_{\Delta,0}]]$ is noetherian, any ideal in it is closed. So all the coefficients of μ with positive exponent that are contained in $J_{\alpha,0}$ can be removed recursively this way: first the smallest j . Therefore we find an element $\mu'' \in J$ such that $\mu''_i \notin J_{\alpha,0}$ and for all $j > 0$ we either have $\mu''_j = 0$ or $\mu''_j \notin J_{\alpha,0}$. Let $0 < l = p^r l'$ ($p \nmid l'$) be the smallest integer with the property that $\mu''_l \neq 0$ and l is divisible by the least possible power of p among these indices. Since I is T_0 -invariant and $\lambda_{\alpha^\vee}(1 + p^t)$ is in T_0 for $t > 1$, we have

$$\begin{aligned} I \ni \lambda_\alpha(1 + p^t)\mu'' - \mu'' &= \sum_{j=0}^{\infty} \mu''_j \left(((X_\alpha + 1)^{1+p^t} - 1)^j - X_\alpha^j \right) = \\ &= \sum_{j=1}^{\infty} \mu''_j \left((X_\alpha + X_\alpha^{p^t} + X_\alpha^{p^t+1})^j - X_\alpha^j \right). \end{aligned}$$

For $j = p^k j'$ with $p \nmid j'$ the lowest degree term of $(X_\alpha + X_\alpha^{p^t} + X_\alpha^{p^t+1})^j - X_\alpha^j$ is $j' X_\alpha^{p^k(j'-1)} X_\alpha^{p^k+t}$. Now for an $l \neq j > 0$ with $\mu''_l \neq 0$ we have either $k = r$ and $j' > l'$ or $k > r$ (by the choice of l). Any case we have $p^k(j'-1) + p^{k+t} > p^r(l'-1) + p^{r+t}$ for any such j as soon as we choose t so that $p^{t+1} - p^t > l' - 1$ (since $p^{k+t} - p^{r+t} \geq p^r(p^{t+1} - p^t)$). With such a choice of t we deduce that $\frac{\lambda_\alpha(1+p^t)\mu'' - \mu''}{X_\alpha^{p^r(l'-1)+p^{r+t}}}$ lies in $J = I \cap \kappa[[N_{\Delta,0}]]$ and has constant term $l' \mu''_l$ that does not lie in $J_{\alpha,0}$. This is a contradiction. \square

Now we claim that $J_{\alpha,0} \subseteq J \cap \kappa[[N_{\Delta \setminus \{\alpha\},0}]]$ and hence $J \cap \kappa[[N_{\Delta \setminus \{\alpha\},0}]]$ is nonzero. For an element $\lambda \in J_{\alpha,0}$ choose an element $\mu \in J$ with $\mu_0 = \lambda$. If $\mu_j \neq 0$ for some $j > 0$ then in view of the Lemma choose $\nu_j \in J$ with $\nu_{j,0} = \mu_j$ and let $\mu' := \mu - X_\alpha^j \nu_j$. By the same recursive argument as in the Lemma we find an element $\mu'' \in J$ with $\mu''_0 = \mu_0 = \lambda$ and $\mu''_j = 0$ for $j > 0$ showing the claim. The statement of the proposition follows from the inductual hypothesis: $(J \cap \kappa[[N_{\Delta \setminus \{\alpha\},0}]])[X_\beta^{-1} \mid \beta \in \Delta \setminus \{\alpha\}]$ is a nonzero T_0 -invariant ideal in $\kappa((N_{\Delta \setminus \{\alpha\},0})$ therefore contains 1. \square

2.2 A functor from smooth B -representations to étale T_+ -modules over $A((N_{\Delta,0}))$

We have the following generalization of Lemma 2.6 in [8].

Proposition 2.3. *Let M be a finitely generated module over $A[[N_{\Delta,0}]] [F_\Delta]$ with a semilinear action of T_0 such that M is admissible and smooth as a module over $A[[N_{\Delta,0}]]$ (ie. the Pontryagin dual M^\vee is finitely generated over $A[[N_{\Delta,0}]]$). Then the module $M^\vee[1/X_\alpha \mid \alpha \in \Delta]$ has naturally the structure of an étale T_+ -module over $A((N_{\Delta,0}))$.*

In order to simplify notation we put $X_\Delta := \prod_{\alpha \in \Delta} X_\alpha$ so that we have $(\cdot)[1/X_\Delta] = (\cdot)[1/X_\alpha \mid \alpha \in \Delta]$.

Proof. By passing to the modules $\varpi^r M / \varpi^{r+1} M$ ($0 \leq r < h$) we may assume without loss of generality that $h = 1$. Let C_α be the cokernel of the map $\kappa[[N_{\Delta,0}]] \otimes_{\varphi_\alpha} M \xrightarrow{1 \otimes F_\alpha} M$. Since M is finitely generated over $\kappa[[N_{\Delta,0}]] [F_\beta \mid \beta \in \Delta]$, C_α is finitely generated over the smaller ring $\kappa[[N_{\Delta,0}]] [F_\beta \mid \beta \in \Delta \setminus \{\alpha\}]$. Let $m_1, \dots, m_r \in C_\alpha$ be the generators. Since M is smooth as a representation of $N_{\alpha,0} \leq N_{\Delta,0}$, so is C_α . Therefore there exists a power X_α^s ($s > 0$) of X_α killing each m_i ($1 \leq i \leq r$). However, X_α is in the centre of $\kappa[[N_{\Delta,0}]] [F_\beta \mid \beta \in \Delta \setminus \{\alpha\}]$ as each F_β ($\beta \neq \alpha$) commutes with X_α . Therefore we have $X_\alpha^s C_\alpha = 0$. In particular, we deduce $C_\alpha^\vee[1/X_\alpha] = 0$. This shows that the map

$$M^\vee[1/X_\Delta] \xrightarrow{(1 \otimes F_\alpha)^\vee[1/X_\Delta]} (\kappa[[N_{\Delta,0}]] \otimes_{\varphi_\alpha, \kappa[[N_{\Delta,0}]]} M)^\vee[1/X_\Delta] \cong \kappa[[N_{\Delta,0}]] \otimes_{\varphi_\alpha, \kappa[[N_{\Delta,0}]]} M^\vee[1/X_\Delta] \quad (1)$$

is injective. Moreover, the generic rank over $\kappa((N_{\Delta,0}))$ of the two sides of (1) equals. Therefore the cokernel of (1) is a finitely generated torsion module over $\kappa((N_{\Delta,0}))$ since M is admissible. Moreover, the global annihilator of this cokernel is T_0 -invariant as the map (1) is T_0 -equivariant. Proposition 2.1 shows that in fact (1) is an isomorphism for each $\alpha \in \Delta$. \square

For a smooth representation π of B_+ over A we can make $\pi^{H_{\Delta,0}}$ a module over $A[[N_{\Delta,0}]] [F_\Delta]$ by the Hecke action $F_\alpha(m) := \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0} t_\alpha^{-1})} u t_\alpha m$. Let us denote by $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ the set of those finitely generated $A[[N_{\Delta,0}]] [F_\Delta]$ -submodules M of $\pi^{H_{\Delta,0}}$ that are stable under the action of T_0 and are admissible as a module over $A[[N_{\Delta,0}]]$. We define

$$D_\Delta^\vee(\pi) := \varprojlim_{M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})} M^\vee[1/X_\Delta].$$

Using Prop. 2.3 this is a projective limit of finitely generated étale T_+ -module over $A((N_{\Delta,0}))$ attached functorially to π : If $f: \pi \rightarrow \pi'$ is a morphism of smooth A -representations of B and M lies in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ then $f(M)$ lies in $\mathcal{M}_\Delta(\pi'^{H_{\Delta,0}})$. Indeed, $f(M)$ is finitely generated over $A[[N_{\Delta,0}]] [F_\Delta]$, stable under the action of T_0 , and admissible as a representation of N_Δ .

2.3 The category $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$

For a submonoid $T_* \leq T_+$ we denote by $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ the category of finitely generated étale T_* -modules over $A((N_{\Delta,0}))$. We regard these objects as left modules over $A((N_{\Delta,0}))$. Further, we denote by $\mathcal{D}^{pro-et}(T_*, A((N_{\Delta,0})))$ the category of projective limits of objects in $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$.

Remark. It is shown in Cor. 3.16 in [29] that any object D in $\mathcal{D}^{et}(T_+, \kappa((N_{\Delta,0})))$ is free as a module over $\kappa((N_{\Delta,0}))$. However, we do not use this fact in the present paper.

Since $\kappa((N_{\Delta,0}))$ is a localization of the local noetherian ring $\kappa[[N_{\Delta,0}]]$ it is also noetherian and has finite global dimension ($\leq |\Delta|$). Moreover, any module over $\kappa((N_{\Delta,0}))$ admits a free resolution of finite length since any projective module over $\kappa[[N_{\Delta,0}]]$ is free. In particular, we may define the (generic) rank of a module M over $\kappa((N_{\Delta,0}))$ as the alternating sum $\text{rk}(M) := \text{rk}_{\kappa((N_{\Delta,0}))}(M) := \sum_{i=0}^{|\Delta|} (-1)^i r_i$ for a free resolution

$$0 \rightarrow F_{|\Delta|} \rightarrow \cdots \rightarrow F_0 \rightarrow M \rightarrow 0$$

with $F_i \cong \kappa((N_{\Delta,0}))^{r_i}$ ($i = 0, \dots, |\Delta|$). This is equal to the dimension of $Q((N_{\Delta,0})) \otimes_{\kappa((N_{\Delta,0}))} M$ where $Q((N_{\Delta,0}))$ denotes the field of fractions of $\kappa((N_{\Delta,0}))$.

For a (left) module D over $A((N_{\Delta,0}))$ we define the generic length of D as $\text{length}_{gen}(D) := \text{length}_{gen, A((N_{\Delta,0}))}(D) := \sum_{i=0}^{h-1} \text{rk}_{\kappa((N_{\Delta,0}))}(\varpi^i D / \varpi^{i+1} D)$. Let $Q(A((N_{\Delta,0})))$ be the localization of $A((N_{\Delta,0}))$ at the prime ideal generated by ϖ . This is an artinian local ring with maximal ideal generated by ϖ and residue field isomorphic to $Q((N_{\Delta,0}))$. We have an isomorphism

$$Q((N_{\Delta,0})) \otimes_{\kappa((N_{\Delta,0}))} (\varpi^i D / \varpi^{i+1} D) \cong \varpi^i Q(A((N_{\Delta,0}))) \otimes_{A((N_{\Delta,0}))} D / \varpi^{i+1} Q(A((N_{\Delta,0}))) \otimes_{A((N_{\Delta,0}))} D.$$

Therefore the generic length of an $A((N_{\Delta,0}))$ -module D equals the length of $Q(A((N_{\Delta,0}))) \otimes_{A((N_{\Delta,0}))} D$. In particular, the generic length is additive on short exact sequences.

Lemma 2.4. *For a finitely generated module D in $A((N_{\Delta,0}))$ and $t \in T_+$ we have $\text{length}_{gen}(D) = \text{length}_{gen}(A((N_{\Delta,0})) \otimes_{\varphi_t} D)$.*

Proof. Note that we have $\varpi^i A((N_{\Delta,0})) \otimes_{\varphi_t} D / \varpi^{i+1} A((N_{\Delta,0})) \otimes_{\varphi_t} D \cong \kappa((N_{\Delta,0})) \otimes_{\varphi_t} (\varpi^i D / \varpi^{i+1})$. So we may assume $A = \kappa$ and $\text{length}_{gen} = \text{rk}$. The statement is clear since $\varphi_t: \kappa((N_{\Delta,0})) \rightarrow \kappa((N_{\Delta,0}))$ is flat, so it takes free resolutions to free resolutions. \square

Proposition 2.5. *If D_2 is an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ and D_1 is a T_+ -stable $A((N_{\Delta,0}))$ -submodule then both D_1 and D_2/D_1 are étale for the inherited action of T_+ , ie. objects in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$. In particular, $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ is an abelian category.*

Proof. It is clear that $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ is an additive category. So it suffices to show the first statement. Put $D_3 := D_2/D_1$ and for a fixed $t \in T_+$ consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A((N_{\Delta,0})) \otimes_{\varphi_t} D_1 & \longrightarrow & A((N_{\Delta,0})) \otimes_{\varphi_t} D_2 & \longrightarrow & A((N_{\Delta,0})) \otimes_{\varphi_t} D_3 \longrightarrow 0 \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 \\ 0 & \longrightarrow & D_1 & \longrightarrow & D_2 & \longrightarrow & D_3 \longrightarrow 0 \end{array}$$

with exact rows. Since f_2 is an isomorphism, we deduce that f_1 is injective and f_3 is surjective. Therefore $\text{Coker}(f_1)$ and $\text{Ker}(f_3)$ have 0 generic length by Lemma 2.4. In particular, $\text{Coker}(f_1)/\varpi\text{Coker}(f_1)$ and $\text{Ker}(f_3)/\varpi\text{Ker}(f_3)$ are finitely generated torsion modules over $\kappa((N_{\Delta,0}))$ both admitting a semilinear action of T_0 . Hence their global annihilator is a nonzero T_0 -invariant ideal in $\kappa((N_{\Delta,0}))$ that contains 1 by Prop. 2.1. We obtain that $\text{Coker}(f_1)/\varpi\text{Coker}(f_1) = \text{Ker}(f_3)/\varpi\text{Ker}(f_3) = 0$ whence we also have $\text{Coker}(f_1) = \text{Ker}(f_3) = 0$ showing that both f_1 and f_3 are isomorphisms. \square

Remark. The above proof actually shows that $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ is an abelian category for any submonoid $T_* \leq T_+$ containing an open subgroup of T_0 .

2.4 A functor to usual (φ, Γ) -modules

Assume in this section that $\Delta \neq \emptyset$ or, equivalently, that $G \neq T$. There exists a $\xi(\mathbb{Z}_p \setminus \{0\})$ -equivariant group homomorphism

$$\ell = \ell_{gen} := \sum_{\alpha \in \Delta} u_\alpha^{-1} : N_{\Delta,0} \twoheadrightarrow \mathbb{Z}_p .$$

We denote by $H_{\ell, \Delta, 0} \leq N_{\Delta,0}$ the kernel of the group homomorphism ℓ and by $H_{\ell,0}$ its preimage in N_0 under the quotient map $N_0 \twoheadrightarrow N_{\Delta,0}$. The restriction of ℓ to $N_{\alpha,0}$ is an isomorphism for all $\alpha \in \Delta$, so ℓ is generic. Moreover, the induced ring homomorphism $\ell : A[[N_{\Delta,0}]] \twoheadrightarrow A[[\mathbb{Z}_p]] \cong A[[X]]$ extends to a surjective ring homomorphism $\ell : A((N_{\Delta,0})) \twoheadrightarrow A((X))$. Its kernel is generated by the elements $X_\alpha - X_\beta$ for $\alpha, \beta \in \Delta$. So we obtain a functor $A((X)) \otimes_{\ell, A((N_{\Delta,0}))} \cdot$ from the category $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ of étale T_+ -modules over $A((N_{\Delta,0}))$ to the category $\mathcal{D}^{et}(\varphi, \Gamma, A((X)))$ of étale (φ, Γ) -modules over $A((X))$.

Proposition 2.6. *The functor $A((X)) \otimes_{\ell, A((N_{\Delta,0}))} \cdot$ from $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ to $\mathcal{D}^{et}(\varphi, \Gamma, A((X)))$ is faithful and exact. In particular, if D is a nonzero étale T_+ -module over $A((N_{\Delta,0}))$ then $D_\ell := A((X)) \otimes_{\ell, A((N_{\Delta,0}))} D$ is nonzero either.*

Proof. At first we prove the exactness. The functor $A((X)) \otimes_{\ell, A((N_{\Delta,0}))} \cdot$ is clearly right exact. We show by induction on $|\Delta|$ that it takes injective maps to injective maps. If $|\Delta| = 1$ then there is nothing to prove. So let $|\Delta| > 1$ and choose $\alpha \neq \beta \in \Delta$. Denote by $T_{+, \alpha=\beta}$ the submonoid in T_+ on which the two characters α and β agree and put $T_{0, \alpha=\beta} := T_0 \cap T_{+, \alpha=\beta}$. We have a functor $A((N_{\Delta,0}))/ (X_\alpha - X_\beta) \otimes_{A((N_{\Delta,0}))} \cdot$ from $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ to the category $\mathcal{D}^{et}(T_{+, \alpha=\beta}, A((N_{\Delta,0}))/ (X_\alpha - X_\beta))$ of étale T_+ -modules over $A((N_{\Delta,0}))/ (X_\alpha - X_\beta)$. Let

$$0 \rightarrow D_1 \rightarrow D_2 \rightarrow D_3 \rightarrow 0$$

be an exact sequence in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$. By induction it suffices to show that the sequence

$$0 \rightarrow D_1 / (X_\alpha - X_\beta) \rightarrow D_2 / (X_\alpha - X_\beta)$$

is exact. Assume that $d_1 + (X_\alpha - X_\beta)D_1$ lies in the kernel of the above map for some $d_1 \in D_1$. Then we have $d_1 \in D_1 \cap (X_\alpha - X_\beta)D_2$ (viewing D_1 as a subobject of D_2). Therefore there exists a $d_2 \in D_2$ such that $d_1 = (X_\alpha - X_\beta)d_2$. Then the image d_3 of d_2 in D_3 is killed by $(X_\alpha - X_\beta)$. Assume that $d_3 \neq 0$. Then there is an integer $0 \leq r < h$ such that $d_3 \in \varpi^r D_3 \setminus \varpi^{r+1} D_3$.

However, $\varpi^r D_3 / \varpi^{r+1} D_3$ is torsion-free as a module over $\kappa((N_{\Delta,0}))$ since the global annihilator of its torsion part would be a T_0 -invariant ideal that does not exist by Prop. 2.1. This is a contradiction as the class of d_3 in $\varpi^r D_3 / \varpi^{r+1} D_3$ is killed by $X_\alpha - X_\beta$. So we conclude that d_2 lies in D_1 whence $d_1 + (X_\alpha - X_\beta)D_1 = 0$.

For the faithfulness let $f: D_1 \rightarrow D_2$ be a nonzero map in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$. By passing to a suitable subquotient $\varpi^r D_2 / \varpi^{r+1} D_2$ we may assume without loss of generality that $A = k$. Since $\mathcal{D}^{et}(T_+, \kappa((N_{\Delta,0})))$ is an abelian category, $f(D_1)$ is a subobject in D_2 . By the above left exactness it suffices to show that $\kappa((X)) \otimes_{\ell, \kappa((N_{\Delta,0}))} f(D_1)$ is nonzero. However, this is clear since $f(D_1)$ is torsionfree as a module over $A((N_{\Delta,0}))$ (again by Prop. 2.1), therefore its localization at the maximal ideal $\text{Ker}(\ell)$ of $\kappa((N_{\Delta,0}))$ is nonzero showing that $\kappa((X)) \otimes_{\ell, \kappa((N_{\Delta,0}))} f(D_1) \neq 0$. \square

Remarks. 1. It is shown in [29] that the functor $A((X)) \otimes_{\ell, A((N_{\Delta,0}))} \cdot$ from $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ to $\mathcal{D}^{et}(\varphi, \Gamma, A((X)))$ corresponds to restriction to the diagonal embedding of $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ into its $|\Delta|$ th direct power on the Galois side. In particular, it is not full if $|\Delta| > 1$.

2. We can extend the functor $A((X)) \otimes_{\ell, A((N_{\Delta,0}))} \cdot$ to $\mathcal{D}^{pro-et}(T_+, A((N_{\Delta,0})))$ the following way. For $\hat{D} = \varprojlim_{i \in I} D_i$ with D_i being an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$, we define

$$A((X)) \hat{\otimes}_{\ell, A((N_{\Delta,0}))} \hat{D} := \varprojlim_{i \in I} A((X)) \otimes_{\ell, A((N_{\Delta,0}))} D_i .$$

Then $A((X)) \hat{\otimes}_{\ell, A((N_{\Delta,0}))} \cdot$ is an exact and faithful functor from $\mathcal{D}^{pro-et}(T_+, A((N_{\Delta,0})))$ to $\mathcal{D}^{pro-et}(\varphi, \Gamma, A((X)))$ since $A((X))$ is an artinian ring therefore the category of pseudo-compact modules over $A((X))$ have exact projective limits. For any smooth representation π of B over A we have a surjective map $D_\xi^\vee(\pi) \rightarrow A((X)) \hat{\otimes}_{\ell, A((N_{\Delta,0}))} D_\Delta^\vee(\pi)$. Whether or not this is always an isomorphism (or even in the case π satisfies certain admissibility conditions) is an open question. We shall see later on that for π in the category SP_A this is true.

Let D be an object in $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ for some submonoid $T_* \leq T_+$. For any set of representatives $J(N_{\Delta,0}/tN_{\Delta,0}t^{-1}) \subset N_{\Delta,0}$ and elements $t \in T_+$, $d \in D$ we can write d uniquely as $d = \sum_{u \in J(N_{\Delta,0}/tN_{\Delta,0}t^{-1})} u\varphi_t(d_{t,u})$. As usual [26] [17] we put $\psi_t(d) := d_{t,1}$ for a subset $J(N_{\Delta,0}/tN_{\Delta,0})$ containing 1. We have $\psi_t(u^{-1}d) = d_{t,u}$ for any $u \in J(N_{\Delta,0}/tN_{\Delta,0}t^{-1})$ and $t \in T_*$. Moreover, we have $\psi_t(\lambda\varphi_t(d)) = \psi_t(\lambda)d$ and $\psi_t(\varphi_t(\lambda)d) = \lambda\psi_t(d)$ for all $d \in D$ and $\lambda \in A((N_{\Delta,0}))$ (here we define $\psi_t(\lambda)$ similarly as above—note that $A((N_{\Delta,0}))$ with the conjugation action of T_* is also an object in $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$). Further, for $t_1, t_2 \in T_*$ we have $\psi_{t_1 t_2} = \psi_{t_2} \circ \psi_{t_1}$. We call this the ψ -action of T_+ on D .

Lemma 2.7. *Let D be an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ and d be in D . Choose the set $1 \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})$ of representatives so that whenever $\ell(u)$ and $\ell(v)$ are in the same coset in $\mathbb{Z}_p/p\mathbb{Z}_p$ for some $u, v \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})$ then actually we have $\ell(u) = \ell(v)$. (This can be done as we have a splitting so that $N_{\Delta,0}$ decomposes as $N_{\Delta,0} \cong \text{Ker}(\ell) \times \mathbb{Z}_p$.) Then for the element $1 \otimes d \in A((X)) \otimes_{\ell, A((N_{\Delta,0}))} D = D_\ell$ we have $\psi_s(1 \otimes d) = 1 \otimes (\sum_{u \in J(\text{Ker}(\ell)/s\text{Ker}(\ell)s^{-1})} \psi_s(u^{-1}d))$.*

Proof. We may decompose d as $d = \sum_{u \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})} u\varphi_s \circ \psi_s(u^{-1}d)$. The image of this equation in D_ℓ reads as

$$1 \otimes d = 1 \otimes \sum_{u \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})} u\varphi_s \circ \psi_s(u^{-1}d) = \sum_{u \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})} \ell(u)\varphi_s(1 \otimes \psi_s(u^{-1}d)) .$$

Now since we have $\text{Ker}(\ell) \cap sN_{\Delta,0}s^{-1} = s\text{Ker}(\ell)s^{-1}$ (as conjugation by s on $\text{Im}(\ell) = \mathbb{Z}_p$ is injective), we obtain that $\ell(u)$ runs through a set of representatives of $\mathbb{Z}_p/p\mathbb{Z}_p$ and each value is taken $|\text{Ker}(\ell)/s\text{Ker}(\ell)s^{-1}|$ -times by our assumption on $J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})$. So we compute

$$1 \otimes d = \sum_{v \in J(\mathbb{Z}_p/p\mathbb{Z}_p)} v \varphi_s \left(\sum_{u \in J(N_{\Delta,0}/sN_{\Delta,0}s^{-1}), \ell(u)=v} 1 \otimes \psi_s(u^{-1}d) \right).$$

Since 1 lies in $J(N_{\Delta,0}/sN_{\Delta,0}s^{-1})$, $1 = \ell(1)$ lies in $J(\mathbb{Z}_p/p\mathbb{Z}_p)$. So we deduce the statement by the uniqueness of decomposition of $1 \otimes d$ as a sum above. \square

2.5 Right exactness of D_{Δ}^{\vee}

Assume $T_* = T_+$ and let D_0 be a finitely generated $A[[N_{\Delta,0}]]$ -submodule in an object D in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ that is stable under the ψ -action of T_+ . Since D_0 is compact in its canonical topology, its Pontryagin dual $D_0^{\vee} = \text{Hom}_A^{cont}(D_0, A)$ is discrete. Moreover, the group $N_{\Delta,0}$ acts on D_0^{\vee} by the formula $um(d) := m(u^{-1}d)$ ($u \in N_{\Delta,0}$, $m \in D_0^{\vee}$, $d \in D_0$). This extends to an action of the Iwasawa algebra $A[[N_{\Delta,0}]]$ by continuity making D_0^{\vee} a discrete left $A[[N_{\Delta,0}]]$ -module. For an element $m \in D_0^{\vee}$ and a simple root $\alpha \in \Delta$ we define $F_{\alpha}(m) \in D_0^{\vee}$ by the formula $F_{\alpha}(m)(d) := m(\psi_{t_{\alpha}}(d))$. Denote by $\#$ the continuous (anti-)involution on the (commutative!) ring $A[[N_{\Delta,0}]]$ that sends group elements $u \in N_{\Delta,0} \subset A[[N_{\Delta,0}]]$ to $u^{\#} := u^{-1}$. We have $\varphi_t(\lambda^{\#}) = \varphi_t(\lambda)^{\#}$ for any $\lambda \in A[[N_{\Delta,0}]]$ and $t \in T_+$ since this is true if $\lambda = u$ in $N_{\Delta,0}$. For $\lambda \in A[[N_{\Delta,0}]]$ and $m \in D_0^{\vee}$ we have $F_{\alpha}(\lambda m) = \varphi_{\alpha}(\lambda)F_{\alpha}(m)$ in D_0^{\vee} since

$$\begin{aligned} F_{\alpha}(\lambda m)(d) &= (\lambda m)(\psi_{t_{\alpha}}(d)) = m(\lambda^{\#}\psi_{t_{\alpha}}(d)) = \\ &= m(\psi_{t_{\alpha}}(\varphi_{t_{\alpha}}(\lambda^{\#})d)) = F_{\alpha}(m)(\varphi_{t_{\alpha}}(\lambda)^{\#}d) = (\varphi_{t_{\alpha}}(\lambda)F_{\alpha}(m))(d). \end{aligned}$$

Moreover, we have $F_{\alpha} \circ F_{\beta} = F_{\beta} \circ F_{\alpha}$ on D_0^{\vee} as $\psi_{t_{\alpha}}$ and $\psi_{t_{\beta}}$ commute on D_0 (T_+ is commutative). So by this action of the operators F_{α} ($\alpha \in \Delta$) we equipped D_0^{\vee} with the structure of a module over the ring $A[[N_{\Delta,0}]][[F_{\Delta}]]$.

On the other hand, for each $t \in T_+$ the ψ -action induces a $A[[N_{\Delta,0}]]$ -module homomorphism

$$\begin{aligned} D_0 &\rightarrow A[[N_{\Delta,0}]] \otimes_{\varphi_t, A[[N_{\Delta,0}]]} D_0 \\ d &\mapsto \sum_{u \in J(N_{\Delta,0}/tN_{\Delta,0}t^{-1})} u \otimes \psi_t(u^{-1}d). \end{aligned} \quad (2)$$

The above map (2) is injective for all $t \in T_+$ (ie. the ψ -action is nondegenerate in the sense of section 4 in [17]) since D_0 is a ψ -invariant submodule of an étale T_+ -module. Therefore in this case $D_0[1/X_{\Delta}]$ is an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$: the map

$$D_0[1/X_{\Delta}] \rightarrow A((N_{\Delta,0})) \otimes_{\varphi_t, A((N_{\Delta,0}))} D_0[1/X_{\Delta}]$$

induced by (2) is a T_0 -equivariant injective map between modules of equal generic length, so it is an isomorphism by Prop. 2.1.

Proposition 2.8. *Let D be an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ and $D_0 \subseteq D$ be a finitely generated $A[[N_{\Delta,0}]]$ -submodule that is stable under the ψ -action of T_+ . Then D_0^{\vee} is a finitely generated module over $A[[N_{\Delta,0}]][[F_{\Delta}]]$ that is admissible as a (smooth) representation of $N_{\Delta,0}$ and admits a semilinear action of T_0 .*

Proof. The case $G = T$ is trivial, so we assume $\Delta \neq \emptyset$.

The admissibility of D_0^\vee follows from the assumption that $(D_0^\vee)^\vee \cong D_0$ is a finitely generated module over $A[[N_{\Delta,0}]]$. Similarly, T_0 acts on D_0 therefore also on D_0^\vee . It remains to show that D_0^\vee is finitely generated as a module over $A[[N_{\Delta,0}]][[F_\Delta]]$. By the above discussion we see that $D_0[1/X_\Delta]$ is a subobject in D (in the category $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$) containing D_0 therefore we may assume without loss of generality that $D = D_0[1/X_\Delta]$ (Prop. 2.5). Denote by $D_{\ell,0}$ the image of D_0 under the surjective map $D \rightarrow D_\ell$ sending $d \in D$ to $1 \otimes d$. By Lemma 2.7 $D_{\ell,0} \subset D_\ell$ is a ψ - and Γ -invariant treillis in the étale (φ, Γ) -module D_ℓ . By Lemma 3.5 in [17] $D_{\ell,0}^\vee$ is a finitely generated module over the ring $A[[X]][[F]]$. Let $m_1, \dots, m_r \in D_{\ell,0}^\vee$ be a finite set of generators. Note that $D_{\ell,0}^\vee$ is contained in the $H_{\ell,\Delta,0}$ -invariant part $(D_0^\vee)^{H_{\ell,\Delta,0}}$ since $H_{\ell,\Delta,0}$ acts trivially on it. (Recall from section 2.4 that $H_{\ell,\Delta,0} \leq N_{\Delta,0}$ is the kernel of the group homomorphism $\ell: N_{\Delta,0} \rightarrow \mathbb{Z}_p$.) Moreover, the action of the ring $A[[X]][[F]]$ comes from the ring homomorphism

$$\begin{aligned} A[[N_{\Delta,0}]][\mathrm{Tr}_{H_{\ell,\Delta,0}/sH_{\ell,\Delta,0}s^{-1}} \circ \prod_{\alpha \in \Delta} F_\alpha] &\twoheadrightarrow A[[X]][[F]] \\ A[[N_{\Delta,0}]] \ni \lambda &\mapsto \ell(\lambda) \\ \mathrm{Tr}_{H_{\ell,\Delta,0}/sH_{\ell,\Delta,0}s^{-1}} \circ \prod_{\alpha \in \Delta} F_\alpha &\mapsto F \end{aligned}$$

where we consider

$$\mathrm{Tr}_{H_{\ell,\Delta,0}/sH_{\ell,\Delta,0}s^{-1}} \circ \prod_{\alpha \in \Delta} F_\alpha = \sum_{u \in J(H_{\ell,\Delta,0}/sH_{\ell,\Delta,0}s^{-1})} u \prod_{\alpha \in \Delta} F_\alpha$$

as an element in $A[[N_{\Delta,0}]][[F_\Delta]]$. Denote by M the $A[[N_{\Delta,0}]][[F_\Delta]]$ -submodule of D_0^\vee generated by the elements $t_0 m_i$ with $t_0 \in T_0$ and $1 \leq i \leq r$. By the discussion above we deduce that $D_{\ell,0}^\vee$ is contained in $M^{H_{\ell,\Delta,0}}$. On the other hand, the orbit of each m_i ($1 \leq i \leq r$) under T_0 is finite. Indeed, if m_i lies in $(D_0^\vee)^{N_{\Delta,0}^{p^n}}$ for some integer n then we also have $t_0 m_i \in (D_0^\vee)^{N_{\Delta,0}^{p^n}}$ since the subgroup $N_{\Delta,0}^{p^n}$ is normalized by T_0 . However, $(D_0^\vee)^{N_{\Delta,0}^{p^n}}$ is finite by the admissibility. In particular, M is finitely generated over $A[[N_{\Delta,0}]][[F_\Delta]]$. Therefore by Proposition 2.3 $M^\vee[1/X_\Delta]$ has the structure of an étale T_+ -module over $A((N_{\Delta,0}))$. The surjective map $D_0 \twoheadrightarrow M^\vee$ induces a surjective morphism $f: D \rightarrow M^\vee[1/X_\Delta]$. The map $D \rightarrow D_\ell$ factors through f since $D_{\ell,0}^\vee$ is contained in M . From Prop. 2.6 we deduce that f is an isomorphism as $D_\ell \xrightarrow{\sim} (M^\vee[1/X_\Delta])_\ell$ by construction. Since D_0 is a subspace in D we obtain that $D_0 \cong M^\vee$ whence $D_0^\vee = M$ is finitely generated over $A[[N_{\Delta,0}]][[F_\Delta]]$. \square

Proposition 2.9. *Let π_1 be a smooth subrepresentation of the smooth A -representation π_2 of B and $M \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})$. Then there exists an $M_1 \subseteq M \cap \pi_1$ such that M_1 lies in $\mathcal{M}_\Delta(\pi_1^{H_{\Delta,0}})$ and $M_1^\vee[1/X] = (M \cap \pi_1)^\vee[1/X]$.*

Proof. The case $G = T$ is trivial, so we assume $\Delta \neq \emptyset$. By the exactness of Pontryagin duality and localization we have an exact sequence

$$0 \rightarrow (M/(M \cap \pi_1))^\vee[1/X_\Delta] \rightarrow M^\vee[1/X_\Delta] \rightarrow (M \cap \pi_1)^\vee[1/X_\Delta] \rightarrow 0.$$

Note that $M/(M \cap \pi_1)$ is finitely generated over $A[[N_{\Delta,0}]][[F_\Delta]]$ and admissible as a representation of $N_{\Delta,0}$ (being a quotient of the finitely generated and admissible M), and it admits a

semilinear action of T_0 . Therefore by Prop. 2.3 and 2.5 all three modules in the above short exact sequence are étale T_+ -modules. So if $M_1 \leq M \cap \pi_1$ is the submodule dual to the image of $(M \cap \pi_1)^\vee$ in $(M \cap \pi_1)^\vee[1/X_\Delta]$ then M_1 is an object in $\mathcal{M}_\Delta(\pi_1^{H_{\Delta,0}})$ by Prop. 2.8. \square

Remark. If we assume further that the map $M^\vee \rightarrow M^\vee[1/X]$ is injective (ie. M^\vee is torsion-free) then we may choose $M_1 = M \cap \pi_1$.

Our main result in this section is the following

Theorem 2.10. *The functor D_Δ^\vee is right exact.*

Proof. The case $G = T$ is trivial, so we assume $\Delta \neq \emptyset$. The proof is now similar to that of Prop. 2.7(ii) in [8], however, we needed Prop. 2.6 in the preparation. Let $0 \rightarrow \pi_1 \rightarrow \pi_2 \rightarrow \pi_3$ be an exact sequence of smooth A -representations of B . Then we have an exact sequence $0 \rightarrow \pi_1^{H_{\Delta,0}} \xrightarrow{f_1} \pi_2^{H_{\Delta,0}} \xrightarrow{f_2} \pi_3^{H_{\Delta,0}}$. Now if M_i is in $\mathcal{M}_\Delta(\pi_i^{H_{\Delta,0}})$ then $f_i(M_i)$ is in $\mathcal{M}_\Delta(\pi_{i+1}^{H_{\Delta,0}})$ ($i = 1, 2$) since the image of a finitely generated module over a ring is again finitely generated and the admissible representations of $N_{\Delta,0}$ form an abelian category. Now let M_2 be in $\mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})$ and put $M_3 := f_2(M_2)$. Then we have $M_3 \in \mathcal{M}_\Delta(\pi_3^{H_{\Delta,0}})$ and by Prop. 2.9 there exists an M_1 in $\mathcal{M}_\Delta(\pi_1^{H_{\Delta,0}})$ such that we have an exact sequence

$$0 \rightarrow M_3^\vee[1/X] \rightarrow M_2^\vee[1/X] \rightarrow M_1^\vee[1/X] \rightarrow 0$$

of objects in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$. Note that for $M_2 \subseteq M_2'$ we also have $M_3 \subseteq M_3' = f_2(M_2')$. Therefore the projective system $(f(M_2)^\vee[1/X])_{M_2 \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})}$ satisfies the Mittag-Leffler property since for $M_2, M_2' \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})$ we also have $M_2 + M_2' \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})$. So by taking the projective limit we obtain an exact sequence

$$0 \rightarrow \varprojlim_{M_2 \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})} f_2(M_2)^\vee[1/X] \rightarrow \varprojlim_{M_2 \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})} M_2^\vee[1/X] \rightarrow \varprojlim_{M_2 \in \mathcal{M}_\Delta(\pi_2^{H_{\Delta,0}})} (M_2 \cap \pi_1)^\vee[1/X] \rightarrow 0$$

In the above short exact sequence the left term is the image of $D_\Delta^\vee(\pi_3)$ in $D_\Delta^\vee(\pi_2)$, the middle term is $D_\Delta^\vee(\pi_2)$, and the term on the right hand side equals $D_\Delta^\vee(\pi_1)$ since all $M_1 \in \mathcal{M}_\Delta(\pi_1^{H_{\Delta,0}})$ can also be viewed as a subspace in $\pi_2^{H_{\Delta,0}}$. \square

3 Compatibility with tensor products and parabolic induction

Our goal in this section is to generalize the results in sections 5–7 in [8] to the functor D_Δ^\vee .

3.1 Tensor products

As in section 5 in [8] let G' be another \mathbb{Q}_p -split reductive group over \mathbb{Q}_p with connected centre and Borel subgroup B' with \mathbb{Q}_p -split torus T' and unipotent radical N' with compact open subgroup N'_0 that we also assume to be totally decomposed. We denote by Φ^+ (resp. by Δ') the set of positive (resp. simple) roots corresponding to B' . For each $\alpha' \in \Phi^+$ we fix isomorphisms $u_{\alpha'} : \mathbb{G}_a \rightarrow N_{\alpha'}$ for the root subgroups $N_{\alpha'} \leq N'$ such that $u_{\alpha'}(\mathbb{Z}_p) = N_{\alpha'} \cap N'_0 =:$

$N'_{\alpha',0}$. Since the centre of G' is connected, there exists a system $(\lambda_{\alpha'^{\vee}})_{\alpha' \in \Delta'} \in X^\vee(T')^{\Delta'}$ of cocharacters with the property $\beta' \circ \lambda_{\alpha'^{\vee}} = 1$ for all $\alpha' \neq \beta' \in \Delta'$ and $\alpha' \circ \lambda_{\alpha'^{\vee}} = \text{id}_{\mathbf{G}_m}$. As in [8] and [25] we put $\xi := \sum_{\alpha' \in \Delta'} \lambda_{\alpha'^{\vee}}$. We define $N'_{\Delta',0} := \prod_{\alpha' \in \Delta'} N'_{\alpha',0}$ arising naturally as a quotient of the group N'_0 . We put $H'_{\Delta',0} := \text{Ker}(N'_0 \rightarrow N'_{\Delta',0})$. Similarly, we consider the rings $A((N'_{\Delta',0}))$ and $A[[N'_{\Delta',0}]] [F_{\Delta'}]$ and consider étale T'_+ -modules over the former forming the categories $\mathcal{D}^{et}(T'_+, A((N'_{\Delta',0})))$ and $\mathcal{D}^{pro-et}(T'_+, A((N'_{\Delta',0})))$ in the usual sense.

We consider the group $G \times G'$ with Borel $B \times B'$ and torus $T \times T'$. We also form the rings $A((N_{\Delta,0} \times N'_{\Delta',0}))$ and $A[[N_{\Delta,0} \times N'_{\Delta',0}]] [F_{\Delta'}]$ and consider étale $T_+ \times T'_+$ -modules over the former ring forming the categories $\mathcal{D}^{et}(T_+ \times T'_+, A((N_{\Delta,0} \times N'_{\Delta',0})))$ and $\mathcal{D}^{pro-et}(T_+ \times T'_+, A((N_{\Delta,0} \times N'_{\Delta',0})))$ in the usual sense.

For finitely generated modules M_0 (resp. M'_0) over the Iwasawa algebra $A[[N_{\Delta,0}]]$ (resp. over $A[[N'_{\Delta',0}]]$) the completed tensor product $M_0 \hat{\otimes}_A M'_0 := (M_0^\vee \otimes_A M'_0{}^\vee)^\vee$ is a finitely generated module over the Iwasawa algebra $A[[N_{\Delta,0} \times N'_{\Delta',0}]]$. It is isomorphic to the projective limit of $M_* \otimes_A M'_*$ where M_* (resp. M'_*) runs through the finite quotients of M_0 (resp. of M'_0).

Now let M (resp. M') be a finitely generated module over $A((N_{\Delta,0}))$ (resp. over $A((N'_{\Delta',0}))$) and choose a finitely generated $A[[N_{\Delta,0}]]$ -submodule $M_0 \subset M$ (resp. $A[[N'_{\Delta',0}]]$ -submodule $M'_0 \subset M'$) such that we have $M = M_0[1/X_\Delta]$ (resp. $M' = M'_0[1/X_{\Delta'}]$) (we call these submodules lattices in M (resp. in M')). We define $M \hat{\otimes}_A M' := (M_0 \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}]$. This is a finitely generated module over $A((N_{\Delta,0} \times N'_{\Delta',0}))$.

Lemma 3.1. *$M \hat{\otimes}_A M'$ does not depend on the choice of M_0 and M'_0 up to a natural isomorphism.*

Proof. Let $M_1 \subset M$ (resp. $M'_1 \subset M'$) be another lattice. It suffices to treat the case when $M'_1 = M'_0$. Assume first that $M_1 = X_\Delta^n M_0$ for some (positive) integer n . Since the multiplication by X_Δ^n on M_0 induces the multiplication by $(X_\Delta^\#)^n$ on the Pontryagin dual, it induces again the multiplication by X_Δ^n on the completed tensor product $M_0 \hat{\otimes}_A M'_0$ which becomes an isomorphism after inverting $X_\Delta X_{\Delta'}$. So the natural inclusion of M_1 in M induces an isomorphism $(M_0 \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}] \cong (M_1 \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}]$.

Now let M_1 be arbitrary. Since we have $M = M_1[1/X_\Delta] = M_0[1/X_\Delta]$, there exist positive integers r_1, r_2 such that $M_1 \subseteq X_\Delta^{r_1} M_0 \subseteq X_\Delta^{r_2} M_1 (\subseteq X_\Delta^{r_1+r_2} M_0)$. By the above case these inclusions induce a sequence of map

$$\begin{aligned} & (M_1 \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}] \xrightarrow{f_1} ((X_\Delta^{r_1} M_0) \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}] \xrightarrow{f_2} \\ & \xrightarrow{f_3} ((X_\Delta^{r_2} M_1) \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}] \xrightarrow{f_4} ((X_\Delta^{r_1+r_2} M_0) \hat{\otimes}_A M'_0)[1/X_\Delta X_{\Delta'}] \end{aligned}$$

such that $f_2 \circ f_1$ and $f_4 \circ f_3$ are isomorphisms. We obtain that f_2 is both injective and surjective therefore an isomorphism. This combined with the above case we obtain statement. The isomorphism is induced by the inclusions $M_1 \cap M_0 \subseteq M_0$ and $M_1 \cap M_0 \subseteq M_1$. \square

Whenever D (resp. D') is an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ (resp. in $\mathcal{D}^{et}(T'_+, A((N'_{\Delta',0})))$) then the completed tensor product $D \hat{\otimes} D'$ can be equipped with the structure of an étale $T_+ \times T'_+$ -module over $A((N_{\Delta,0} \times N'_{\Delta',0}))$ the following way. Choose an $A[[N_{\Delta,0}]]$ -lattice D_0 (resp. an $A[[N'_{\Delta',0}]]$ -lattice D'_0) in D (resp. in D') and an element $t \in T_+$. The action of φ_t on D provides us with an isomorphism

$$1 \otimes \varphi_t: \varphi_t^* D_0 \rightarrow A[[N_{\Delta,0}]] \varphi_t(D_0). \quad (3)$$

(Here we put $\varphi_t^* D_0 := A[[N_{\Delta,0}]] \otimes_{\varphi_t} D_0$ as in [17].) Further, we compute

$$\begin{aligned} (\varphi_t^* D_0) \hat{\otimes}_A D'_0 &= ((\varphi_t^* D_0)^\vee \otimes_A D_0^{\vee})^\vee \cong (A[[N_{\Delta,0}]] \otimes_{\varphi_t, A[[N_{\Delta,0}]]} D_0^\vee \otimes_A D_0^{\vee})^\vee \cong \\ &\cong A[[N_{\Delta,0}]] \otimes_{\varphi_t, A[[N_{\Delta,0}]]} (D_0^\vee \otimes_A D_0^{\vee})^\vee \cong A[[N_{\Delta,0}]] \otimes_{\varphi_t, A[[N_{\Delta,0}]]} (D_0 \hat{\otimes}_A D'_0) \cong \\ &\cong A[[N_{\Delta,0} \times N'_{\Delta',0}]] \otimes_{\varphi_t, A[[N_{\Delta,0} \times N'_{\Delta',0}]]} (D_0 \hat{\otimes}_A D'_0) \end{aligned}$$

as $t \in T_+$ acts trivially on $N'_{\Delta',0}$. By the above identification the isomorphism (3) induces an isomorphism

$$1 \otimes (\varphi_t \otimes 1): A[[N_{\Delta,0} \times N'_{\Delta',0}]] \otimes_{\varphi_t, A[[N_{\Delta,0} \times N'_{\Delta',0}]]} (D_0 \hat{\otimes}_A D'_0) \rightarrow (A[[N_{\Delta,0}]] \varphi_t(D_0)) \hat{\otimes}_A D'_0 .$$

Inverting $X_\Delta X_{\Delta'}$ and noting that $A[[N_{\Delta,0}]] \varphi_t(D_0)$ is a lattice in D (as D is an étale T_+ -module) we obtain an isomorphism

$$1 \otimes (\varphi_t \otimes 1): A((N_{\Delta,0} \times N'_{\Delta',0})) \otimes_{\varphi_t, A((N_{\Delta,0} \times N'_{\Delta',0}))} (D \hat{\otimes}_A D') \rightarrow D \hat{\otimes}_A D' .$$

By symmetry, we obtain a similar isomorphism for any $t' \in T'_+$. This equips $D \hat{\otimes}_A D'$ with the structure of an étale $T_+ \times T'_+$ -module over $A((N_{\Delta,0} \times N'_{\Delta',0}))$. Now if $D = \varprojlim_{i \in I} D_i$ (resp. $D' = \varprojlim_{i' \in I'} D'_{i'}$) is an object in $\mathcal{D}^{pro-et}(T_+, A((N_{\Delta,0})))$ (resp. in $\mathcal{D}^{pro-et}(T'_+, A((N'_{\Delta',0})))$) then we may form the completed tensor product $D \hat{\otimes}_A D' := \varprojlim_{i \in I, i' \in I'} D_i \hat{\otimes}_A D'_{i'}$. This is an object in $\mathcal{D}^{pro-et}(T_+ \times T'_+, A((N_{\Delta,0} \times N'_{\Delta',0})))$.

Proposition 3.2. *Let π (resp. π') be a smooth representation of B (resp. of B') over κ . Then we have an isomorphism*

$$D_{\Delta \cup \Delta'}^\vee(\pi \otimes_\kappa \pi') \cong D_\Delta^\vee(\pi) \hat{\otimes}_\kappa D_{\Delta'}^\vee(\pi')$$

in $\mathcal{D}^{pro-et}(T_+ \times T'_+, \kappa((N_{\Delta,0} \times N'_{\Delta',0})))$.

Proof. We proceed in 3 steps. *Step 1.* We show that we have

$$(\pi \otimes_\kappa \pi')^{H_{\Delta,0} \times H'_{\Delta',0}} = \pi^{H_{\Delta,0}} \otimes_\kappa \pi^{H'_{\Delta',0}} . \quad (4)$$

Since κ is a field, \otimes_κ is exact, therefore the right hand side of (4) is can be viewed as a subspace in $\pi \otimes_\kappa \pi'$. The inclusion \supseteq in (4) is therefore clear. Let now $\sum_{i=1}^r m_i \otimes m'_i$ be an arbitrary element in the left hand side of (4). Since κ is a field, we may assume that the elements $m'_1, \dots, m'_r \in \pi'$ are linearly independent over κ . Now for any $h \in H_{\Delta,0}$ we have

$$\sum_{i=1}^r m_i \otimes m'_i = h \left(\sum_{i=1}^r m_i \otimes m'_i \right) = \sum_{i=1}^r (hm_i) \otimes m'_i .$$

Since the elements m'_i ($1 \leq i \leq r$) are linearly independent, we deduce that $m_i = hm_i$ for all $1 \leq i \leq r$ showing that $\sum_{i=1}^r m_i \otimes m'_i$ lies in $\pi^{H_{\Delta,0}} \otimes_\kappa \pi'$. Now we may rewrite $\sum_{i=1}^r m_i \otimes m'_i$ so that the $m_i \in \pi^{H_{\Delta,0}}$ ($1 \leq i \leq r$) are linearly independent (possibly losing the assumption that m'_i ($1 \leq i \leq r$) are linearly independent). By the same process with $H'_{\Delta',0}$ we deduce (4).

Step 2. Now let M (resp. M') be an object in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ (resp. in $\mathcal{M}_{\Delta'}(\pi'^{H'_{\Delta',0}})$). Then $M \otimes_\kappa M'$ is T_0 -invariant and admissible as a representation of $N_{\Delta,0} \times N'_{\Delta',0}$ since for an open

subgroup $N_{\Delta,*} \leq N_{\Delta,0}$ (resp. $N'_{\Delta',*} \leq N'_{\Delta',0}$) the product $N_{\Delta,*} \times N'_{\Delta',*}$ is open in $N_{\Delta,0} \times N'_{\Delta',0}$ and the κ -vectorspace $(M \otimes_{\kappa} M')^{N_{\Delta,*} \times N'_{\Delta',*}} = M^{N_{\Delta,*}} \otimes_{\kappa} M'^{N'_{\Delta',*}}$ is finite dimensional (using again the argument above). Moreover, choose generators m_1, \dots, m_r (resp. $m'_1, \dots, m'_{r'}$) of M (resp. of M') as a module over $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ (resp. over $\kappa[[N'_{\Delta',0}]] [F_{\Delta'}]$). Then the elements $m_i \otimes m'_j$ ($1 \leq i \leq r$, $1 \leq j \leq r'$) generate $M \otimes_{\kappa} M'$ as a module over $\kappa[[N_{\Delta,0} \times N'_{\Delta',0}]] [F_{\Delta \cup \Delta'}]$ since $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ and $\kappa[[N'_{\Delta',0}]] [F_{\Delta'}]$ are subrings of this ring and any element in $M \otimes_{\kappa} M'$ can be written as a sum of elements of the form $m \otimes m'$ where $m \in M$ and $m' \in M'$. So $M \otimes_{\kappa} M'$ is an object in $\mathcal{M}_{\Delta \cup \Delta'}((\pi \otimes_{\kappa} \pi')^{H_{\Delta,0} \times H'_{\Delta',0}})$.

Step 3: We show that the elements of the form $M \otimes_{\kappa} M'$ for M (resp. M') in $\mathcal{M}_{\Delta}(\pi^{H_{\Delta,0}})$ (resp. in $\mathcal{M}_{\Delta'}(\pi'^{H'_{\Delta',0}})$) are cofinal in $\mathcal{M}_{\Delta \cup \Delta'}((\pi \otimes_{\kappa} \pi')^{H_{\Delta,0} \times H'_{\Delta',0}})$. Let M'' be in $\mathcal{M}_{\Delta \cup \Delta'}((\pi \otimes_{\kappa} \pi')^{H_{\Delta,0} \times H'_{\Delta',0}})$ with a finite set $m''_1, \dots, m''_{r''}$ of generators as a module over $\kappa[[N_{\Delta,0} \times N'_{\Delta',0}]] [F_{\Delta \cup \Delta'}]$ and by (4) write each of the m''_i ($1 \leq i \leq r''$) as a sum of elementary tensors of the form $m \otimes m'$ with $m \in \pi^{H_{\Delta,0}}$ and $m' \in \pi'^{H'_{\Delta',0}}$. Let U (resp. U') be the set of $m \in \pi^{H_{\Delta,0}}$ (resp. of $m' \in \pi'^{H'_{\Delta',0}}$) appearing in these elementary tensors. These are finite sets. We may assume without loss of generality that the elements of the set U' are linearly independent and each $m' \in U'$ appears only once in each m''_i ($1 \leq i \leq r''$). Moreover, for each $m' \in U'$ let $U(m') \subseteq U$ be the subset of those $m \in U$ such that $m \otimes m'$ appears as an elementary tensor in one of the m''_i ($1 \leq i \leq r''$). We then have $U = \bigcup_{m' \in U'} U(m')$.

Lemma 3.3. *The $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -submodule of $\pi^{H_{\Delta,0}}$ generated by the elements of U is admissible as a representation of $N_{\Delta,0}$.*

Proof. Since the subring $\kappa[[N_{\Delta,0}]] [F_{\Delta}] \leq \kappa[[N_{\Delta,0} \times N'_{\Delta',0}]] [F_{\Delta \cup \Delta'}]$ acts via the first component on the tensor product $M \otimes_{\kappa} M'$, any element m_* in $\sum_{i=1}^{r''} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m''_i$ can be uniquely written as a sum $m_* = \sum_{m' \in U'} m_*(m') \otimes m'$ for some elements $m_*(m') \in \sum_{m \in U(m')} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m$. Since the representation π' of B' is smooth and U' is a finite set, there exists an open subgroup $N'_{\Delta',*} \leq N'_{\Delta',0}$ stabilizing all the elements m' in U' . Now for an open subgroup $N_{\Delta,*} \leq N_{\Delta,0}$, an element $m_* = \sum_{m' \in U'} m_*(m') \otimes m'$ in $\sum_{i=1}^{r''} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m''_i$ is stabilized by $N_{\Delta,*} \times N'_{\Delta',*}$ if and only if each $m_*(m')$ ($m' \in U'$) is stabilized by $N_{\Delta,*}$. Since $M \otimes_{\kappa} M'$ is admissible as a representation of $N_{\Delta,0} \times N'_{\Delta',0}$, we deduce that

$$\left(\sum_{i=1}^{r''} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m''_i \right)^{N_{\Delta,*}}$$

is finite dimensional for any open subgroup $N_{\Delta,*} \leq N_{\Delta,0}$. In particular, $\sum_{i=1}^{r''} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m''_i$ is admissible as a representation of $N_{\Delta,0}$. Now for each $m'_0 \in U'$ consider the projection

$$\begin{aligned} \Pi_{m'_0} : \sum_{m' \in U'} \pi^{H_{\Delta,0}} \otimes m' &\rightarrow \pi^{H_{\Delta,0}} \\ \sum_{m'_* \in U'} m(m') \otimes m' &\mapsto m(m'_0) . \end{aligned}$$

$\Pi_{m'_0}$ is $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -linear. Moreover, we have

$$\Pi_{m'_0} \left(\sum_{i=1}^{r''} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m''_i \right) = \sum_{m \in U(m'_0)} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m .$$

In particular, $\sum_{m \in U(m'_0)} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m$ is admissible as a representation of $N_{\Delta,0}$ being the image of an admissible representation. We deduce the statement from the equality $U = \bigcup_{m' \in U'} U(m')$ noting that the finite sum of admissible representations is also admissible. \square

We choose a basis $U_* \subset U$ of the κ -vectorspace generated by $m \in U$. Now by possibly grouping together some elementary tensors in m''_i ($1 \leq i \leq r''$), we may write each element m''_i as a finite sum of elementary tensors $m''_i = \sum_{m \in U_*} m \otimes m'_i(m)$. By using again the above Lemma in the symmetric situation we deduce that the $\kappa[[N'_{\Delta',0}]] [F_{\Delta'}]$ -submodule of $\pi'^{H'_{\Delta',0}}$ generated by $U'_* := \{m'_i(m) \mid 1 \leq i \leq r'', m \in U_*\}$ is admissible as a representation of $N'_{\Delta',0}$. Moreover, M'' is clearly contained in $M \otimes_{\kappa} M'$ where M (resp. M') is the $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -submodule (resp. $\kappa[[N'_{\Delta',0}]] [F_{\Delta'}]$ -submodule) of $\pi^{H_{\Delta,0}}$ (resp. of $\pi'^{H'_{\Delta',0}}$) generated by U_* (resp. by U'_*). Now if we replace U_* and U'_* by their finite T_0 -orbit (resp. T'_0 -orbit) we may assume that M and M' are stable under the action of T_0 (resp. of T'_0). This finishes the proof of Step 3.

The statement follows from the cofinality of the $\kappa[[N_{\Delta,0} \times N'_{\Delta',0}]] [F_{\Delta \times \Delta'}]$ -submodules of the form $M \otimes_{\kappa} M'$ for $M \in \mathcal{M}_{\Delta}(\pi^{H_{\Delta,0}})$ and $M' \in \mathcal{M}_{\Delta'}(\pi'^{H'_{\Delta',0}})$ in $\mathcal{M}_{\Delta \cup \Delta'}((\pi \otimes_{\kappa} \pi')^{H_{\Delta,0} \times H'_{\Delta',0}})$. \square

Remark. The above proof seems to only work in the case of κ -representations, but not for A -representations. However, it might be possible to deduce the general case (or at least the case when the representations are on free A -modules) from the above Proposition. Note that in case $A = \kappa$ the above statement is slightly stronger than Prop. 5.5 in [8] since we do not assume the property Ad (or its analogue in this situation) in Def. 2.2 in [8] that all the finitely generated $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -submodules (resp. $\kappa[[N'_{\Delta',0}]] [F_{\Delta'}]$ -submodules) of $\pi^{H_{\Delta,0}}$ (resp. of $\pi'^{H'_{\Delta',0}}$) are admissible as representations of $N_{\Delta,0}$ (resp. of $N'_{\Delta',0}$) (see also Remark 5.6(ii) of [8]).

3.2 Parabolic induction

As in [8] let $P \leq G$ be a parabolic subgroup containing B with Levi component L_P and unipotent radical $N_P \leq N$. Denote by P^- the opposite parabolic and by N_{P^-} its unipotent radical. The group L_P is also a \mathbb{Q}_p -split reductive group and we fix its Borel subgroup $B_{L_P} := B \cap L_P$ with unipotent radical $N_{L_P} := N \cap L_P$. We denote by Φ_P the roots of the pair (L_P, T) and by $\Phi_P^+ \subseteq \Phi_P$ the subset of positive roots with respect to B_{L_P} and by $\Delta_P \subseteq \Phi_P^+$ the set of simple roots. Since the centre $Z(G)$ is connected, the \mathbb{Z} -module $X(T)/(\bigoplus_{\alpha \in \Delta} \mathbb{Z}\alpha)$ is torsion-free. Therefore $X(T)/(\bigoplus_{\alpha \in \Delta_P} \mathbb{Z}\alpha)$ is torsion-free, too, because $\bigoplus_{\alpha \in \Delta_P} \mathbb{Z}\alpha$ is a direct summand in $\bigoplus_{\alpha \in \Delta} \mathbb{Z}\alpha$.

We adapt the constructions of the previous sections to the group L_P : We denote by $N_{L_P,0} := N_0 \cap N_{L_P}$ and $N_{\Delta_P,0} := \prod_{\alpha \in \Delta_P} N_{\alpha,0}$. We regard $N_{\Delta_P,0}$ as a direct summand in $N_{\Delta,0}$ and also as a quotient of the group $N_{L_P,0}$. We denote by $H_{\Delta_P,0}$ the kernel of the quotient map $N_{L_P,0} \twoheadrightarrow N_{\Delta_P,0}$.

As in [8] denote by $W := N_G(T)/T$ (resp. by $W_P := N_{L_P}(T)/T$) the Weyl group of G (resp. of L_P) and by $w_0 \in W$ the element of maximal length. We have a canonical system

$$K_P := \{w \in W \mid w^{-1}(\Phi_P^+) \subseteq \Phi^+\}$$

of representatives (the Kostant representatives) of the right cosets $W_P \backslash W$. We have a generalized Bruhat decomposition

$$G = \coprod_{w \in K_P} P^- w B = \coprod_{w \in K_P} P^- w N .$$

Now let π_P be a smooth representation of L_P over A . We regard π_P as a representation of P^- via the quotient map $P^- \rightarrow L_P$. Then the parabolically induced representation $\text{Ind}_{P^-}^G \pi_P$ admits [27] (see also [15] §4.3) a filtration by B -subrepresentations whose graded pieces are contained in

$$\mathcal{C}_w(\pi_P) := c - \text{Ind}_{P^-}^{P^- w N} \pi_P$$

for $w \in K_P$ where $c - \text{Ind}$ stands for the space of locally constant functions with compact support modulo P^- . B acts on $\mathcal{C}_w(\pi_P)$ by right translations. Moreover, the first graded piece equals $\mathcal{C}_1(\pi_P)$.

Lemma 3.4. *Let $\pi' \leq \mathcal{C}_w(\pi_P)$ be any B -subrepresentation for some $w \in K_P \setminus \{1\}$. Then we have $D_\Delta^\vee(\pi') = 0$.*

Proof. By the right exactness (and contravariance) of D_Δ^\vee (Thm. 2.10) it suffices to treat the case $\pi' = \mathcal{C}_w(\pi_P)$. This follows from Prop. 6.2 in [8] using that $A((X)) \otimes_{\ell, A((N_{\Delta, 0}))} \cdot$ is faithful (Prop. 2.6) and that we have a surjective map from $D_\xi^\vee(\pi')$ to $A((X)) \otimes_{\ell, A((N_{\Delta, 0}))} D_\Delta^\vee(\pi')$ by the Remark after Prop. 2.6. \square

Theorem 3.5. *Let π_P be a smooth locally admissible representation of L_P over A which we view by inflation as a representation of P^- . We have an isomorphism*

$$D_\Delta^\vee(\text{Ind}_{P^-}^G \pi_P) \cong A((N_{\Delta, 0})) \hat{\otimes}_{A((N_{\Delta_P, 0}))} D_{\Delta_P}^\vee(\pi_P)$$

in the category $\mathcal{D}^{\text{pro-}et}(T_+, A((N_{\Delta, 0})))$.

Remark. Here $\hat{\otimes}$ stands for taking tensor products on finitely generated quotients and then inverse limits.

Proof. By Lemma 3.4 and Thm. 2.10 it suffices to show the isomorphism $D_\Delta^\vee(\mathcal{C}_1(\pi_P)) \cong A((N_{\Delta, 0})) \hat{\otimes}_{A((N_{\Delta_P, 0}))} D_{\Delta_P}^\vee(\pi_P)$.

The map $\mathcal{C}_1(\pi_P) \rightarrow c - \text{Ind}_{N_{L_P}}^N \pi_P$ sending a function $f: P^- N \rightarrow \pi_P$ to its restriction to N is a B -equivariant isomorphism. So we make this identification. Denote by $\mathcal{C}_{1,0}(\pi_P)$ the B_+ -subrepresentation of $\mathcal{C}_1(\pi_P)$ of functions supported on $N_{L_P} N_0$ and let M be arbitrary in $\mathcal{M}_\Delta(\mathcal{C}_1(\pi_P)^{H_{\Delta, 0}})$. So we have a short exact sequence

$$0 \rightarrow (M/M \cap \mathcal{C}_{1,0}(\pi_P))^\vee [1/X_\Delta] \rightarrow M^\vee [1/X_\Delta] \rightarrow (M \cap \mathcal{C}_{1,0}(\pi_P))^\vee [1/X_\Delta] \rightarrow 0$$

in $\mathcal{D}^{et}(T_+, A((N_{\Delta, 0})))$. By Lemma 6.6 in [8] and Prop. 2.6 we deduce that the left hand side $(M/M \cap \mathcal{C}_{1,0}(\pi_P))^\vee [1/X_\Delta]$ vanishes. On the other hand, by Prop. 2.8 we find as in the proof of Prop. 2.9 a finitely generated $A[[N_{\Delta, 0}]] [F_\Delta]$ -submodule $M_1 \leq M \cap \mathcal{C}_{1,0}(\pi_P)$ that is T_0 -stable and admissible as a representation of $N_{\Delta, 0}$ such that we have $M_1^\vee [1/X_\Delta] = (M \cap \mathcal{C}_{1,0}(\pi_P))^\vee [1/X_\Delta]$. Therefore we obtain an isomorphism $D_\Delta^\vee(\mathcal{C}_1(\pi_P)) \cong D_{\Delta}^\vee(\mathcal{C}_{1,0}(\pi_P))$.

Now we identify $\mathcal{C}_{1,0}(\pi_P)$ with $\text{Ind}_{N_{LP,0}}^{N_0} \pi_P$. On the latter $t \in T_+$ acts via the formula

$$(tf)(v) = \begin{cases} 0 & \text{if } v \in N_0 \setminus N_{LP,0}tN_0t^{-1}; \\ u_1t \cdot f(v_1) & \text{if } v = u_1tv_1t^{-1} \in N_{LP,0}tN_0t^{-1} \end{cases} = \\ = \sum_{u \in J(N_{LP,0}/tN_{LP,0}t^{-1})} ut \cdot f(t^{-1}u^{-1}vt) \quad (5)$$

by extending in (5) the functions $f \in \text{Ind}_{N_{LP,0}}^{N_0} \pi_P$ to N with 0 outside N_0 .

Lemma 3.6. *We have a B_0 -equivariant identification $\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}} \cong \text{Ind}_{N_{\Delta P,0}}^{N_{\Delta,0}} \pi_P^{H_{\Delta P,0}}$.*

Proof. Let $f: N_0 \rightarrow \pi_P$ be an $H_{\Delta,0}$ -invariant function in $\mathcal{C}_{1,0}(\pi_P)$. Now for any $v \in N_0$ and $h \in H_{\Delta,0}$ we have $f(v) = (hf)(v) = f(vh)$. Hence, since $H_{\Delta,0}$ is normal in N_0 , we deduce $f(h^{-1}vh) = f(v \cdot v^{-1}h^{-1}vh) = f(v)$. Now if we have $h \in H_{\Delta P,0} \leq H_{\Delta,0}$ and $v \in N_0$ then we compute $f(v) = f(h \cdot h^{-1}vh) = h \cdot f(h^{-1}vh) = h \cdot f(v)$. So we obtain that the image of f lies in $\pi_P^{H_{\Delta P,0}}$. Using again that f is $H_{\Delta,0}$ -invariant the function $\tilde{f}: N_{\Delta,0} \rightarrow \pi_P^{H_{\Delta P,0}}$, $\tilde{f}(vH_{\Delta,0}) := f(v)$ is well-defined. Vica versa, if $\tilde{f}: N_{\Delta,0} \rightarrow \pi_P^{H_{\Delta P,0}}$ is a $N_{\Delta P,0}$ -invariant map then we may lift it to a map $f: N_{LP,0} \setminus N_0 / H_{\Delta,0}$. Therefore the map $f \mapsto \tilde{f}$ is a B_0 -equivariant bijection. \square

Lemma 3.7. *Let M be an object in $\mathcal{M}_{\Delta}(\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}})$. Then the set $M_* := \{f(1) \in \pi_P^{H_{\Delta P,0}} \mid f \in M\}$ is an object in $\mathcal{M}_{\Delta P}(\pi_P^{H_{\Delta P,0}})$.*

Proof. We proceed in 3 steps.

Step 1: We show that M_* is an $A[[N_{\Delta P,0}]] [F_{\Delta P}]$ -submodule. At first note that there is an alternative description of M_* as $M_* = \{m \in \pi_P^{H_{\Delta P,0}} \mid \exists f \in M, v \in N_0: f(v) = m\}$ since for any $v \in N_0$ and $f \in M$ we have $(vf)(1) = f(v)$. Let t be in T_+ and f be in $\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}}$ arbitrary and compute

$$F_t(f)(v) = \sum_{u \in J(H_{\Delta,0}/tH_{\Delta,0}t^{-1})} (utf)(v) = \sum_{u \in J(H_{\Delta,0}/tH_{\Delta,0}t^{-1})} (tf)(vu) = \\ = \sum_{u \in J(H_{\Delta,0}/tH_{\Delta,0}t^{-1})} \sum_{u' \in J(N_{LP,0}/tN_{LP,0}t^{-1})} u't \cdot f(t^{-1}u'^{-1}vut) \quad (6)$$

using (5). Set $J := J(H_{\Delta,0}/tH_{\Delta,0}t^{-1})$ and $J' := J(N_{LP,0}/tN_{LP,0}t^{-1})$. Now consider the bipartite graph on the set $J \amalg J'$ in which there is an edge between $u \in J$ and $u' \in J'$ if and only if $t^{-1}u'^{-1}vut$ lies in N_0 . If a pair (u, u') is not an edge then we have $f(t^{-1}u'^{-1}vut) = 0$ as f is supported on N_0 . Note that the degree of each vertex in this graph is at most 1. Whenever v lies in $N_{LP}tN_0t^{-1}H_{\Delta,0}$ then we do have a (non-unique) $u'_0 \in J'$ and a unique $u(u'_0) \in J$ such that $u'_0{}^{-1}vu(u'_0)$ lies in tN_0t^{-1} . Moreover, the class of u'_0 in $N_{\Delta P,0}/tN_{\Delta P,0}t^{-1}$ is unique since $H_{\Delta,0}$ lies in the kernel of the quotient map $N_0 \twoheadrightarrow N_{\Delta,0}$ and the value $f(t^{-1}u'^{-1}vu(u')t)$ only depends on the class of u' in $N_{\Delta P,0}$ for each $u' \in J'$. Since (6) does not depend on the choice

of J' , we may choose $J' := J(N_{L_P,0}/tN_{L_P,0}t^{-1}H_{\Delta_P,0})J(H_{\Delta_P,0}/tH_{\Delta_P,0}t^{-1})$ and deduce

$$\begin{aligned} F_t(f)(v) &= \sum_{\omega \in J(H_{\Delta_P,0}/tH_{\Delta_P,0}t^{-1})} \sum_{u \in J} \sum_{u' \in J(N_{L_P,0}/tN_{L_P,0}t^{-1}H_{\Delta_P,0})} u'\omega t \cdot f(t^{-1}\omega^{-1}u'^{-1}vut) = \\ &= \sum_{u \in J} \sum_{u' \in J(N_{L_P,0}/tN_{\Delta_P,0}t^{-1}H_{\Delta_P,0})} u'F_t(f(t^{-1}u'^{-1}vut)) \end{aligned} \quad (7)$$

so that for each $v \in N_0$ there is at most one pair $(u, u') \in J \times J(N_{L_P,0}/tN_{L_P,0}t^{-1}H_{\Delta_P,0})$ for which $f(t^{-1}u'^{-1}vut) \neq 0$. In particular, we have $F_t(f)(1) = F_t(f(1))$ whence M_* is an $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$ -submodule.

Step 2. We show that M_* is finitely generated over $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$. Now let f_1, \dots, f_r be a finite set of generators of M as a module over $A[[N_{\Delta,0}]] [F_{\Delta}]$. Since each $f_i: N_0 \rightarrow \pi_P^{H_{\Delta_P,0}}$ is continuous, N_0 is compact, $\pi_P^{H_{\Delta_P,0}}$ is discrete, the set $U_0 := \{f_i(v) \mid v \in N_0, 1 \leq i \leq r\}$ is finite. Moreover, since π_P is locally admissible and t_α lies in the centre of L_P for all $\alpha \in \Delta \setminus \Delta_P$, the orbit of the elements in U_0 under the action of $\prod_{\alpha \in \Delta \setminus \Delta_P} t_\alpha^{\mathbb{Z}}$ is also finite. Therefore the union $U := \prod_{\alpha \in \Delta \setminus \Delta_P} t_\alpha^{\mathbb{Z}} U_0$ is also finite. Since f_1, \dots, f_r generates M , we may write any function $f \in M$ as a finite sum of functions of the form $vF_t(f_i)$ for $v \in N_0$, $t \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$, and $1 \leq i \leq r$. We decompose $t = t_1 t_2$ for $t_1 \in \prod_{\alpha \in \Delta_P} t_\alpha^{\mathbb{N}}$ and $t_2 \in \prod_{\alpha \in \Delta \setminus \Delta_P} t_\alpha^{\mathbb{N}}$. By (7) we obtain that

$$(vF_t(f_i))(1) = F_t(f_i)(v) = \sum_{u \in J} \sum_{u' \in J(N_{L_P,0}/tN_{L_P,0}t^{-1}H_{\Delta_P,0})} u'F_{t_1}(F_{t_2}(f_i(t^{-1}u'^{-1}vut)))$$

showing that U generates M_* as a module over $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$ since $F_{t_2}(f_i(t^{-1}u'^{-1}vut))$ lies in U .

Step 3: We show that M_* is admissible as a representation of N_{Δ_P} . For each generator $m \in U$ of M_* choose a function $f_m \in M$ with $f_m(1) = m$. We put $V := \{f_m \in M \mid m \in U\}$. Let N_* be an open subgroup of $N_{\Delta,0}$ stabilizing all the functions in V and put $N_{\Delta \setminus \Delta_P,*} := N_* \cap N_{\Delta \setminus \Delta_P,0}$ where $N_{\Delta \setminus \Delta_P,0} := \prod_{\alpha \in \Delta \setminus \Delta_P} N_{\alpha,0}$. Let t be in $\prod_{\alpha \in \Delta_P} t_\alpha^{\mathbb{N}}$ and v be in $N_{\Delta_P,0}$ be arbitrary. Using $F_t(f)(1) = F_t(f(1))$ which follows from (7) and that every function $f \in \text{Ind}_{N_{\Delta_P,0}}^{N_0} \pi_P$ satisfies $f(v) = v \cdot f(1)$ by definition, we obtain

$$(vF_t(f_m))(1) = F_t(f_m)(v) = v \cdot F_t(f_m)(1) = v \cdot F_t(f_m(1)) = v \cdot F_t(m). \quad (8)$$

This shows that the map

$$M_1 := \sum_{m \in U} A[[N_{\Delta_P,0}]] [F_{\Delta_P}] f_m \ni f \mapsto f(1) \in M_*$$

is a surjective $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$ -module homomorphism. Moreover, both $N_{\Delta_P,0}$ and $\prod_{\alpha \in \Delta_P} t_\alpha^{\mathbb{Z}}$ commute with $N_{\Delta \setminus \Delta_P,0}$, therefore any element in M_1 is $N_{\Delta \setminus \Delta_P,*}$ -invariant. Therefore M_1 is admissible as a representation of $N_{\Delta_P,0}$ since for any open subgroup $N_{\Delta_P,*} \leq N_{\Delta_P,0}$ we have

$$M_1^{N_{\Delta_P,*}} = M_1^{N_{\Delta_P,*} \times N_{\Delta \setminus \Delta_P,*}} \subset M^{N_{\Delta_P,*} \times N_{\Delta \setminus \Delta_P,*}}$$

which is finite. In particular, M_* is also admissible as it arises as a quotient of M_1 .

Finally, it is clear that M_* is invariant under the action of T_0 hence belongs to $\mathcal{M}_{\Delta_P}(\pi_P^{H_{\Delta_P,0}})$. \square

Now let M' be arbitrary in $\mathcal{M}_{\Delta_P}(\pi_P)$ and define $\widetilde{M}' := \{f \in \mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}} \mid f(v) \in M' \text{ for all } v \in N_0\} \cong \text{Ind}_{N_{\Delta_P,0}}^{N_{\Delta,0}} M'$.

Lemma 3.8. \widetilde{M}' lies in $\mathcal{M}_{\Delta}(\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}})$. For any M in $\mathcal{M}_{\Delta}(\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}})$ we have $M \subseteq \widetilde{M}_*$.

Proof. By the definition of M_* and \widetilde{M}_* , M is contained in \widetilde{M}_* .

Let M' be arbitrary in $\mathcal{M}_{\Delta_P}(\pi_P^{H_{\Delta_P,0}})$. By (7), \widetilde{M}' is an $A[[N_{\Delta,0}]] [F_{\Delta}]$ -submodule of $\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}}$. Choose a finite set $m_1, \dots, m_r \in M'$ of generators as a module over $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$. For each $1 \leq i \leq r$ let $f_i \in \mathcal{C}_{1,0}(\pi_P)$ be the constant function $f_i(v) := m_i$ for all $v \in N_0$. Since f_i is constant, it is $H_{\Delta,0}$ -invariant therefore lies in \widetilde{M}' . We claim that the functions f_1, \dots, f_r generate \widetilde{M}' as a module over $A[[N_{\Delta,0}]] [F_{\Delta}]$. Let f be any function in \widetilde{M}' . By Lemma 3.6, we may view f as a function on $N_{\Delta,0}$ that is determined by its values on the subgroup $N_{\Delta \setminus \Delta_P,0}$. Moreover, since the restriction of f to $N_{\Delta \setminus \Delta_P,0}$ is continuous, it is constant on the cosets of subgroup $tN_{\Delta \setminus \Delta_P,0}t^{-1}$ for some $t \in \prod_{\alpha \in \Delta \setminus \Delta_P} t_{\alpha}^{\mathbb{N}}$ as these subgroups form a system of neighbourhoods of 1 in $N_{\Delta \setminus \Delta_P,0}$. So it suffices to show that for each coset $v_0 t N_{\Delta \setminus \Delta_P,0} t^{-1}$ ($v_0 \in N_{\Delta \setminus \Delta_P,0}$) and m in M' the function

$$f_{v_0,t,m}: N_0 \rightarrow \pi_P^{H_{\Delta_P,0}}$$

$$v \mapsto \begin{cases} u \cdot m & \text{if } v \in uv_0 t N_{\Delta_P,0} t^{-1} H_{\Delta,0} \text{ for some } u \in N_{L_P,0}; \\ 0 & \text{otherwise,} \end{cases}$$

lies in $\sum_{i=1}^r A[[N_{\Delta,0}]] [F_{\Delta}] f_i$. Since m_1, \dots, m_r generate M' , we may write m as a finite sum $\sum_{i=1}^r \lambda_i m_i$ for some $\lambda_i \in A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$ ($1 \leq i \leq r$). We have $\sum_{i=1}^r \lambda_i f_i(1) = m$ by (8). On the other hand, each $\lambda_i f_i$ ($1 \leq i \leq r$) is constant on $N_{\Delta \setminus \Delta_P,0}$ since $N_{\Delta \setminus \Delta_P,0}$ commutes with all the elements in $A[[N_{\Delta_P,0}]] [F_{\Delta_P}]$ and f_i is constant on $N_{\Delta \setminus \Delta_P,0}$. Using (7) we deduce $f_{v_0,t,m} = \sum_{i=1}^r v_0 F_t \lambda_i f_i$ since for $t \in \prod_{\alpha \in \Delta \setminus \Delta_P} t_{\alpha}^{\mathbb{N}}$ we have $N_{L_P,0} = t N_{L_P,0} t^{-1} H_{\Delta_P,0}$.

It is clear that \widetilde{M}' is T_0 -stable. Moreover, we have $\widetilde{M}'^{\vee} = A[[N_{\Delta,0}]] \otimes_{A[[N_{\Delta_P,0}]]} M'^{\vee}$. In particular, \widetilde{M}'^{\vee} is finitely generated over $A[[N_{\Delta,0}]]$ since M' is admissible as a representation of $N_{\Delta_P,0}$. \square

So the elements of the form \widetilde{M}' ($M' \in \mathcal{M}_{\Delta_P}(\pi_P)$) are cofinal in $\mathcal{M}_{\Delta}(\mathcal{C}_{1,0}(\pi_P)^{H_{\Delta,0}})$. For these we have $\widetilde{M}'[1/X_{\Delta}] \cong A((N_{\Delta,0})) \otimes_{A((N_{\Delta_P,0}))} M'[1/X_{\Delta_P}]$. We finish the proof of Thm. 3.5 by taking the projective limit. \square

Corollary 3.9. For a character $\chi: T \rightarrow A^{\times}$, the étale T_+ -module $D_{\Delta}^{\vee}(\text{Ind}_B^G \chi) \cong A((N_{\Delta,0})) \otimes_A \chi$ is free of rank 1 on which T_+ acts via the twist by χ .

Question 1. Let π be a supercuspidal representation of G . Is the converse of Theorem 3.5 true in the sense that—assuming $D_{\Delta}^{\vee}(\pi) \neq 0$ —there does not exist a proper subset $I \subsetneq \Delta$ and an étale T_+ -module D over $A((N_{I,0}))$ such that we have

$$D_{\Delta}^{\vee}(\pi) \cong A((N_{\Delta,0})) \hat{\otimes}_{A((N_{I,0}))} D ?$$

3.3 Exactness of D_Δ^\vee on the category SP_A

We start this section with the following abstract Lemmata that will be needed several times in the sequel.

Lemma 3.10. *Let R be an arbitrary ring and $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a short exact sequence of R -modules.*

(i) *If M_1 and M_3 are both finitely presented then so is M_2 .*

(ii) *If M_3 is finitely presented and M_2 is finitely generated then M_1 is finitely generated.*

Proof. This should be classical, but here is a proof: (i) We have a presentation $\bigoplus_{j=1}^{r_l} R d_j^{(l)} \xrightarrow{f_l} \bigoplus_{i=1}^{k_l} R e_i^{(l)} \rightarrow M_l \rightarrow 0$ sending the free generator $e_i^{(l)}$ to some elements $m_i^{(l)} \in M_l$ ($1 \leq i \leq k_l$, $l = 1, 3$) by assumption. We regard M_1 as a submodule of M_2 and lift the elements $m_i^{(3)}$ to $\widetilde{m_i^{(3)}} \in M_2$ ($1 \leq i \leq k_3$). So we have a surjective map

$$\begin{aligned} \Psi: \bigoplus_{i=1}^{k_1} R e_i^{(1)} \oplus \bigoplus_{i'=1}^{k_3} R e_{i'}^{(3)} &\twoheadrightarrow M_2 \\ e_i^{(1)} &\mapsto m_i^{(1)} \\ e_{i'}^{(3)} &\mapsto \widetilde{m_{i'}^{(3)}}. \end{aligned}$$

Moreover, $\text{Ker}(\Psi) \cap \bigoplus_{i=1}^{k_1} R e_i^{(1)} = \text{Im}(f_1)$ and $\text{Ker}(\Psi) / (\text{Ker}(\Psi) \cap \bigoplus_{i=1}^{k_1} R e_i^{(1)}) = \text{Im}(f_3)$ are finitely generated whence so is $\text{Ker}(\Psi)$.

(ii) Let $U \subseteq M_2$ be a finite set of generators containing a set $V \subseteq U$ of lifts of the generators of M_3 arising in the finite presentation of M_3 . Since the image \overline{V} of V in M_3 generates M_3 , we may assume without loss of generality that $U \setminus V$ is contained in M_1 . The natural surjection $\bigoplus_{\overline{v} \in \overline{V}} R e_{\overline{v}} \rightarrow M_3$ sending the formal generator $e_{\overline{v}}$ to \overline{v} has finitely generated kernel M_0 . Moreover, $\sum_{v \in V} \lambda v$ lies in M_1 if and only if $\sum_{v \in V} \lambda v e_{\overline{v}}$ lies in M_0 . This shows that $\sum_{v \in V} R v \cap M_1$ is a quotient of M_0 —in particular, finitely generated. Finally, M_1 is generated by $U \setminus V$ and $\sum_{v \in V} R v \cap M_1$ therefore it is finitely generated. \square

Lemma 3.11. *Let R be an arbitrary ring and $R \leq S = R[s_1, \dots, s_k]$ be a ring extension that is finitely generated as an R -algebra by elements $s_1, \dots, s_k \in S$ commuting with each other and satisfying $s_i R \subseteq R s_i$. If M is an S -module that is finitely presented as an R -module then M is also finitely presented as an S -module.*

Proof. Choose a finite presentation $f: F := \bigoplus_{j=1}^n R e_j \twoheadrightarrow M$ sending e_j to some element $m_j \in M$ ($1 \leq j \leq n$). Consider the map $\text{id} \otimes f: S \otimes_R F = \bigoplus_{j=1}^n S e_j \twoheadrightarrow M$ sending e_j to m_j . Since f is surjective, for each pair (i, j) ($1 \leq i \leq k$, $1 \leq j \leq n$) there exists an element $r_{i,j} \in F$ such that $f(r_{i,j}) = (\text{id} \otimes f)(s_i e_j)$. Let I be the S -submodule of $S \otimes_R F$ generated by $S \otimes_R \text{Ker}(f)$ and $\{s_i e_j - 1 \otimes r_{i,j} \mid 1 \leq i \leq k, 1 \leq j \leq n\}$. It is clear that $I \subseteq \text{Ker}(\text{id} \otimes f)$. On the other hand, let $\sum_{j=1}^n t_j e_j$ be in $\text{Ker}(\text{id} \otimes f)$ ($t_j \in S$, $1 \leq j \leq n$). We may write each t_j as a polynomial in the s_i 's with coefficients in R ($1 \leq i \leq k$, $1 \leq j \leq n$) such that the s_i 's are on the right. Using the relations $s_i e_j - 1 \otimes r_{i,j}$ we find that $\sum_{j=1}^n t_j e_j$ is congruent to an element in $1 \otimes F$ modulo I . So we deduce that $\sum_{j=1}^n t_j e_j$ lies in I so that $I = \text{Ker}(\text{id} \otimes f)$ is finitely generated over S . \square

For a subset $\Delta' \subset \Delta$ we introduce the following notations $N_{\Delta',0} := \prod_{\alpha \in \Delta'} N_{\alpha,0}$ (as a direct summand of the group $N_{\Delta,0}$) and $[F_{\Delta'}] := [F_{\alpha} \mid \alpha \in \Delta']$.

Lemma 3.12. *Let Δ_0 and Δ' be subsets in Δ (with $\Delta_1 := \Delta \setminus \Delta_0$ and $\Delta'' := \Delta \setminus \Delta'$) and let $N_{\Delta_1,*}$ be an open subgroup of $N_{\Delta_1,0}$ such that we have $tN_{\Delta_1,*}t^{-1} \subset N_{\Delta_1,*}$ for all $t \in \prod_{\alpha \in \Delta'} t_{\alpha}^{\mathbb{N}}$. Further, let $\chi: T \rightarrow \kappa^{\times}$ be a continuous character and consider the space $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ of continuous functions as an $N_{\Delta,0}$ -representation with the following action of the operators F_{α} ($\alpha \in \Delta$): For $\alpha \in \Delta'$ we let F_{α} act by zero. For $\alpha \in \Delta''$, $v \in N_{\Delta,0}$ and $f \in \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ we put*

$$F_{\alpha}(f)(v) := \sum_{u \in J((N_{\Delta,0} \cap t_{\alpha}^{-1} N_{\Delta_1,*} t_{\alpha}) / N_{\Delta_1,*})} \chi(t_{\alpha}) f(t_{\alpha}^{-1} v t_{\alpha} u). \quad (9)$$

Then any finitely generated $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -submodule $M \leq \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,}}$ is finitely presented and has finite length. Moreover, if we further assume $\Delta' \subseteq \Delta_1$ then the $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -module $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ is cyclic. If $\Delta' \subseteq \Delta_1$ and $N_{\Delta_1,*} = N_{\Delta_1,0}$ then $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ is simple (ie. has no nontrivial submodules).*

Proof. Step 1: We assume $\Delta' \subseteq \Delta_1$ and show in this case that $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,}}$ is generated by the characteristic function $\mathbb{1} := \mathbb{1}_{N_{\Delta_1,*} \times N_{\Delta_0,0}}$. Note that any function f in $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ is constant on the right cosets of $N_{\Delta_1,*} \times s_0^r N_{\Delta_0,0} s_0^{-r}$ for r large enough depending on f (here we put $s_0 := \prod_{\alpha \in \Delta_0} t_{\alpha}$). Moreover, the function $\chi(s_0^{-r}) u F_{s_0^r}(\mathbb{1})$ is the characteristic function of the right coset $u N_{\Delta_1,*} \times s_0^r N_{\Delta_0,0} s_0^{-r}$ therefore f can be written as a finite κ -linear combination of elements of the form $u F_{s_0^r}(\mathbb{1})$.*

Step 2: We assume $\Delta' \subseteq \Delta_1$ and $N_{\Delta_1,} = N_{\Delta_1,0}$ and show that $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ is simple. Let $0 \neq M \leq \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ be an arbitrary submodule. Then M contains a nonzero fixed point under the action of the pro- p group $N_{\Delta,0}$ therefore also the function $\mathbb{1}_{N_{\Delta,0}}$. By our assumption $\mathbb{1}_{N_{\Delta,0}} = \mathbb{1}_{N_{\Delta_1,*} \times N_{\Delta_0,0}}$ generates $\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$ so we have $M = \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}}$.*

Step 3: We still assume $\Delta' \subseteq \Delta_1$ and $N_{\Delta_1,} = N_{\Delta_1,0}$. We claim that the annihilator left-ideal $I := \text{Ann}(\mathbb{1}_{N_{\Delta,0}}) \triangleleft \kappa[[N_{\Delta,0}]] [F_{\Delta}]$ of $\mathbb{1}_{N_{\Delta,0}}$ is generated by the finite set*

$$U := \{X_{\alpha}, X_{\alpha'}, \chi(t_{\alpha}) - \sum_{u \in J(N_{\Delta_0,0}/t_{\alpha} N_{\Delta_0,0} t_{\alpha}^{-1})} u F_{\alpha}, F_{\alpha'} \mid \alpha' \in \Delta', \alpha \in \Delta''\}.$$

The containment $U \subset I$ is clear. Let $\sum_{i=1}^r \lambda_i F_{t_i} \in I$ be arbitrary ($\lambda_i \in \kappa[[N_{\Delta,0}]]$, $t_i \in \prod_{\alpha \in \Delta} t_{\alpha}^{\mathbb{N}}$, $i = 1, \dots, r$). We may omit the terms $\lambda_i F_{t_i}$ with t_i divisible by $t_{\alpha'}$ for some $\alpha' \in \Delta'$ as those are contained in the left ideal generated by U . Note that for a common multiple $t \in \prod_{\alpha \in \Delta''} t_{\alpha}^{\mathbb{N}}$ there exists a $\lambda \in \kappa[[N_{\Delta,0}]]$ such that we have

$$\sum_{i=1}^r \lambda_i F_{t_i} \equiv \lambda F_t \pmod{\sum_{m \in U} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m}.$$

Indeed, if $t = t_i t'_i$ then we have

$$F_{t_i} \equiv \chi^w(t'_i)^{-1} \sum_{u \in J(N_{\Delta_0,0}/t'_i N_{\Delta_0,0} t'^{-1})} t_i u t_i^{-1} F_t \pmod{\sum_{m \in U} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m}.$$

Further, for $\alpha \in \Delta_1 \cap \Delta''$ we have $t_{\alpha} N_{\Delta_0,0} t_{\alpha}^{-1} = N_{\Delta_0,0}$ whence $\chi(t_{\alpha}) - F_{\alpha}$ lies in U . Therefore we may assume without loss of generality that t lies in $\prod_{\alpha \in \Delta_0} t_{\alpha}^{\mathbb{N}}$ whence $J(N_{\Delta_0,0}/t N_{\Delta_0,0} t^{-1})$

is a set of representatives for the cosets $N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1}$, too. Therefore we may write $\lambda = \sum_{u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})} u\varphi_t(\lambda_{u,t})$ with $\lambda_{u,t} \in \kappa[[N_{\Delta,0}]]$ whence we deduce

$$\begin{aligned} \lambda F_t &= \sum_{u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})} u\varphi_t(\lambda_{u,t})F_t = \sum_{u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})} uF_t\lambda_{u,t} \equiv \\ &\equiv \sum_{u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})} c_{u,t}uF_t \pmod{\sum_{m \in U} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m} \end{aligned}$$

where $c_{u,t} \in \kappa$ is the constant term of $\lambda_{u,t}$ ($u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})$). Now the function $0 = \sum_{u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})} c_{u,t}uF_t(\mathbb{1}_{N_{\Delta_0,0}})$ is constant $c_{u,t}$ on the coset $utN_{\Delta_0,0}t^{-1}$ implying $c_{u,t} = 0$ for each $u \in J(N_{\Delta_0,0}/tN_{\Delta_0,0}t^{-1})$. We obtain $I = \sum_{m \in U} \kappa[[N_{\Delta,0}]] [F_{\Delta}] m$ as claimed.

Step 4: We assume $N_{\Delta_1,*} = N_{\Delta_1,0}$, but drop the assumption that $\Delta' \subseteq \Delta_1$. Choose a finite set U of generators of M as a module over $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$. By assumption $F_{\alpha'}$ ($\alpha' \in \Delta'$) acts trivially on M . On the other hand, F_{α} acts by $\chi(t_{\alpha})$ for each $\alpha \in \Delta_1 \cap \Delta''$. Therefore U generates M as a module over $\kappa[[N_{\Delta,0}]] [F_{\Delta_0 \cap \Delta''}]$, too, and by Lemma 3.11 it suffices to show that it is finitely presented and has finite length as such. Moreover, $N_{\Delta_1,0}$ also acts trivially on M and lies in the centre of the ring $\kappa[[N_{\Delta,0}]] [F_{\Delta_0 \cap \Delta''}]$. So U generates M as a module over $\kappa[[N_{\Delta_0,0}]] [F_{\Delta_0 \cap \Delta''}]$ and by Lemma 3.11 it suffices to show that it is finitely presented and has finite length as such. There exists a subgroup $N_{\Delta_0 \cap \Delta',*} \leq N_{\Delta_0 \cap \Delta',0}$ stabilizing all the elements in U . On the other hand, the subalgebra $\kappa[[N_{\Delta_0 \cap \Delta',0}]]$ lies in the centre of $\kappa[[N_{\Delta_0,0}]] [F_{\Delta'}]$ therefore $N_{\Delta_0 \cap \Delta',*}$ acts trivially on the whole M . Hence the finite orbit $U_1 = N_{\Delta_0 \cap \Delta',0}U$ of U generates M as a module over $\kappa[[N_{\Delta_0 \cap \Delta',0}]] [F_{\Delta_0 \cap \Delta''}]$ and by Lemma 3.11 we are reduced to showing that M is finitely presented and has finite length as such. The elements in M may be regarded as functions on $N_{\Delta_0,0}/N_{\Delta_0 \cap \Delta',*} = N_{\Delta_0 \cap \Delta'',0} \times (N_{\Delta_0 \cap \Delta',0}/N_{\Delta_0 \cap \Delta',*})$. Therefore M is a submodule of

$$\bigoplus_{v \in N_{\Delta_0 \cap \Delta',0}/N_{\Delta_0 \cap \Delta',*}} \mathcal{C}(N_{\Delta_0 \cap \Delta'',0}, \kappa) .$$

Each direct summand above is simple and finitely presented as a module over $\kappa[[N_{\Delta_0 \cap \Delta''}]] [F_{\Delta_0 \cap \Delta''}]$ by Steps 2 and 3 therefore M is also finitely presented by Lemma 3.10 and has finite length.

Step 5: no assumptions. We write $N_{\Delta,0}/N_{\Delta_1,*}$ as a direct product of $N_{\Delta_0,0}$ and the finite p -group $N_{\Delta_1,0}/N_{\Delta_1,*}$ and let J be the Jacobson radical of the group ring $\kappa[N_{\Delta_1,0}/N_{\Delta_1,*}]$. Note that $M^i := J^i \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} \cap M$ is a $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ -submodule of M for all $i \geq 0$. By Lemma 3.10 it suffices to show that each graded piece M^i/M^{i+1} is finitely presented and has finite length ($i \geq 0$). M/M^1 is a finitely generated submodule of

$$\mathcal{C}/\mathcal{C}^1 := \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} / J\mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} \cong \mathcal{C}(N_{\Delta_0,0}, \kappa)$$

on which $\kappa[[N_{\Delta,0}]] [F_{\Delta}]$ acts via its quotient $\kappa[[N_{\Delta_0,0}]] [F_{\Delta}]$. Moreover, all $\alpha' \in \Delta'$ act by zero on $\mathcal{C}/\mathcal{C}^1$. On the other hand, the classes of functions $f \in \mathcal{C}$ supported on $N_{\Delta_1,*} \times N_{\Delta_0,0}$ generate $\mathcal{C}/\mathcal{C}^1$. On such an f each F_{α} ($\alpha \in \Delta''$) acts by the formula $F_{\alpha}(f)(v) = \chi(t_{\alpha})f(t_{\alpha}^{-1}vt_{\alpha})$ since in the sum (9) all the other terms vanish. Therefore M/M^1 is finitely presented and has finite length by Step 4. In particular, M^1 is finitely generated by Lemma 3.10.

Now for $i \geq 1$ we have an identification

$$\mathcal{C}^i/\mathcal{C}^{i+1} := J^i \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} / J^{i+1} \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} \cong J^i/J^{i+1} \otimes_{\kappa} \mathcal{C}/\mathcal{C}^1 .$$

Now J^i/J^{i+1} is generated over κ by the elements $\prod_{\alpha \in \Delta_1} (n_\alpha - 1)^{k_\alpha}$ with $\sum_{\alpha \in \Delta_1} k_\alpha = i$ and n_α topological generator in $N_{\alpha,0}$. On a generator $\prod_{\alpha \in \Delta_1} (n_\alpha - 1)^{k_\alpha}$ the operator F_α ($\alpha \in \Delta_1$) acts by 0 if $k_\alpha \neq 0$ otherwise by 1. Using Step 4 we deduce by induction on i that M^i/M^{i+1} is finitely presented and of finite length (whence M^{i+1} is finitely generated). The statement follows using Lemma 3.10. \square

Following [8] we denote by SP_A the category of smooth G -representations over A consisting of finite length representations whose Jordan-Hölder constituents are subquotients of principal series. Let $\chi: T \rightarrow A^\times$ be a continuous character. The principal series representation $\text{Ind}_B^G \chi$ admits a filtration by B -subrepresentations whose graded pieces are $\mathcal{C}_w(\chi) = c - \text{Ind}_{B^-}^{B^-wN} \chi \cong c - \text{Ind}_{w^{-1}B^-w \cap N}^N \chi^w$ for $w \in W = N_G(T)/T$ where χ^w is the character given by the formula $\chi^w(t) = \chi(wtw^{-1})$. We set $\mathcal{C}_{w,0}(\chi) := \text{Ind}_{B^-}^{B^-wN_0} \chi$ which is a generating B_+ -subrepresentation in $\mathcal{C}_w(\chi)$.

Assume now that $A = \kappa$. We introduce the filtration (indexed by $t \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$)

$$\text{Fil}^t := \text{Fil}^t \mathcal{C}_w(\chi) := \text{Ind}_{w^{-1}B^-w \cap N}^{(w^{-1}B^-w \cap N)t^{-1}N_0t} \chi^w$$

by B_+ -subrepresentations. We have $\text{Fil}^1 = \mathcal{C}_{w,0}(\chi)$. As a representation of N_0 , Fil^t can be written as a direct sum

$$\text{Fil}^t = \bigoplus_{N_{t,v} \in D_{t,w}} \mathcal{C}_{t,v}.$$

where $N_{t,v} := (w^{-1}B^-w \cap t^{-1}N_0t)vN_0$ is the double coset of $v \in t^{-1}N_0t$, $D_{t,w} := (w^{-1}B^-w \cap t^{-1}N_0t) \setminus (t^{-1}N_0t)/N_0$ is the set of double cosets, and $\mathcal{C}_{t,v} := \text{Ind}_{w^{-1}B^-w \cap N}^{N_{t,v}} \chi^w$. Now each $t_* \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ acts on the set $D_{t,w}$ via conjugation: $t_* \cdot N_{t,v} := N_{t,t_*vt_*^{-1}}$. Indeed, this definition does not depend on the choice v of the representative in $N_{t,v}$ since we have $t_*N_0t_*^{-1} \subseteq N_0$ and $t_*(w^{-1}B^-w)t_*^{-1} = w^{-1}B^-w$.

Lemma 3.13. *Assume that we have $t_* \cdot N_{t,v} = N_{t,v}$ for some $t, t_* \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ and $v \in t^{-1}N_0t$. Then there exists a representative v_* in the double coset $N_{t,v}$ such that $t_*v_*t_*^{-1} = v_*$. Moreover, we have $t_\alpha \cdot N_{t,v} = N_{t,v}$ for each $\alpha \in \Delta$ with $\alpha(t_*) \neq 1$.*

Proof. We have $N_{t,v} = N_{t,t_*vt_*^{-1}}$. By induction on i we find that $N_{t,v} = N_{t,t_*^i vt_*^{-i}}$ for all $i \geq 1$. Since $\beta(t_*)$ is a nonnegative power of p for each $\beta \in \Phi^+$, the limit $v_* := \lim_i t_*^i vt_*^{-i}$ exists in N and satisfies $t_*v_*t_*^{-1} = v_*$ and $N_{t,v} = N_{t,v_*}$. Now we can write v_* uniquely in the form $v_* = \prod_{\beta \in \Phi^+} n_\beta$ with $n_\beta \in N_\beta$. So we have $v_* = t_*v_*t_*^{-1} = \prod_{\beta \in \Phi^+} n_\beta^{\beta(t_*)}$. So for each $\beta \in \Phi^+$ we have $n_\beta = 1$ or $\beta(t_*) = 1$. Now if $\alpha\Delta$ is a simple root with $\alpha(t_*) \neq 1$ then we have $n_\beta = 1$ for all $\beta \in \Phi^+$ with $\beta \circ \lambda_{\alpha^\vee} \neq 1$, or equivalently, for all $\beta \in \Phi^+$ with $\beta(t_\alpha) \neq 1$. In particular, we deduce $t_\alpha v_* t_\alpha^{-1} = v_*$ whence $t_\alpha \cdot N_{t,v} = N_{t,v}$. \square

We order the set $D_{t,w}$ partially by putting $N_{t,v_1} \leq N_{t,v_2}$ if there exists an element $t' \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ such that $t' \cdot N_{t,v_2} = N_{t,v_1}$.

Lemma 3.14. *The ordering \leq on $D_{t,w}$ is transitive, reflexive, and antisymmetric.*

Proof. The transitivity and reflexivity are clear ($\prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ is a monoid with $1 \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$). Assume now that $t' \cdot N_{t,v_1} = N_{t,v_2}$ and $t'' \cdot N_{t,v_2} = N_{t,v_1}$. Put $t_* := t't''$. By Lemma 3.13 we deduce $N_{t,v_1} = t'' \cdot N_{t,v_2} = N_{t,v_2}$ as t'' can be written as a product of t_α 's with $\alpha(t_*) \neq 1$. \square

So we can refine the ordering \leq on $D_{t,w}$ to a total ordering of $D_{t,w}$ giving a filtration

$$\text{Fil}^{t,v} := \bigoplus_{N_{t,v} \geq N_{t,v'} \in (w^{-1}B^-w \cap t^{-1}N_0t) \setminus (t^{-1}N_0t)/N_0} \mathcal{C}_{t,v'}$$

on Fil^t by finitely many $N_0 \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ -subrepresentations. The graded piece

$$\text{gr}^{t,v}(\mathcal{C}_w) := \text{Fil}^{t,v} / \bigcup_{N_{t,v'} < N_{t,v}} \text{Fil}^{t,v'}$$

is isomorphic to $\mathcal{C}_{t,v}$ as a representation of N_0 .

Lemma 3.15. *Any finitely generated $\kappa[[N_{\Delta,0}]][[F_\Delta]$ -submodule of $\text{gr}^{t,v}(\mathcal{C}_w)^{H_{\Delta,0}}$ is finitely presented.*

Proof. Put $\Delta_1 := \Delta \cap w^{-1}(\Phi^-)$ and $\Delta_0 := \Delta \setminus w^{-1}(\Phi^-)$. $H_{\Delta,0}$ is normal in N_0 with commutative quotient $N_{\Delta,0}$ such that the image of

$$N_0(t, v) := v^{-1}(w^{-1}B^-w \cap t^{-1}N_0t)v \cap N_0$$

in $N_{\Delta,0}$ is an open subgroup $N_{\Delta_1,*}$ in $N_{\Delta_1,0}$. Hence we have an identification

$$\text{gr}^{t,v}(\mathcal{C}_w)^{H_{\Delta,0}} \cong \text{Ind}_{w^{-1}B^-w \cap t^{-1}N_0t}^{(w^{-1}B^-w \cap t^{-1}N_0t)vN_0/H_{\Delta,0}} \chi^w \cong \mathcal{C}(N_{\Delta,0}, \kappa)^{N_{\Delta_1,*}} \quad (10)$$

as a representation of $N_{\Delta,0}$.

Now we describe the action of each F_α ($\alpha \in \Delta$) on $\text{gr}^{t,v}(\mathcal{C}_w)^{H_{\Delta,0}}$. Let $\alpha \in \Delta$ be arbitrary. If we have $N_{t,v} > N_{t,t_\alpha v t_\alpha^{-1}}$ then F_α acts by 0 on $\text{gr}^{t,v}(\mathcal{C}_w)^{H_{\Delta,0}}$ since even t_α acts by 0 on $\text{gr}^{t,v}(\mathcal{C}_w)$. On the other hand, if $N_{t,v} = N_{t,t_\alpha v t_\alpha^{-1}}$ then by Lemma 3.13 we may assume without loss of generality that the representative v of the double coset $(w^{-1}B^-w \cap t^{-1}N_0t)vN_0$ is chosen so that we have $t_\alpha v t_\alpha^{-1} = v$. Therefore we compute

$$\begin{aligned} F_\alpha(f)(vv_1) &= \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})} (t_\alpha f)(vv_1u) = \\ &= \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})} \chi^w(t_\alpha) f(t_\alpha^{-1}vv_1ut_\alpha) = \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})} \chi^w(t_\alpha) f(vt_\alpha^{-1}v_1ut_\alpha) \end{aligned} \quad (11)$$

for any $v_1 \in N_0$. Now if v_1 does not lie in $N_0(t, v)t_\alpha N_0 t_\alpha^{-1} H_{\Delta,0}$ then all the terms in (11) are zero whence $F_\alpha(f)(v_1) = 0$. However, if $v_1 = n_0 t_\alpha v_2 t_\alpha^{-1} u_0$ for some $n_0 \in N_0(t, v)$, $v_2 \in N_0$, $u_0 \in H_{\Delta,0}$ then we may replace $J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})$ with $J' := u_0 J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})$ and assume without loss of generality that $u_0 = 1$. Moreover, since $t_\alpha v_2 t_\alpha^{-1}$ normalizes both $H_{\Delta,0}$ and $t_\alpha H_{\Delta,0} t_\alpha^{-1}$, $J'' := t_\alpha v_2 t_\alpha^{-1} J' t_\alpha v_2^{-1} t_\alpha^{-1}$ is also a set of representatives of $H_{\Delta,0}/t_\alpha H_{\Delta,0} t_\alpha^{-1}$.

Therefore we compute

$$\begin{aligned}
F_\alpha(f)(vv_1) &= \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})} \chi^w(t_\alpha) f(vt_\alpha^{-1}v_1ut_\alpha) = \\
&= \sum_{u \in J(H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1})} \chi^w(t_\alpha) f(vt_\alpha^{-1}n_0t_\alpha v_2t_\alpha^{-1}u_0ut_\alpha) = \\
&= \sum_{u'' \in J''} \chi^w(t_\alpha) f(vt_\alpha^{-1}n_0u''t_\alpha v_2) = \sum_{\substack{v_2 \in N_0(t,v) \setminus N_0/H_{\Delta,0} \\ v_1 \in N_0(t,v)t_\alpha v_2 t_\alpha^{-1} H_{\Delta,0}}} \sum_{u'' \in N_0(t,v)t_\alpha H_{\Delta,0}t_\alpha^{-1} \cap J''} \chi^w(t_\alpha) f(vv_2) \\
&= \sum_{\substack{v_2 \in N_0(t,v) \setminus N_0/H_{\Delta,0} \\ v_1 \in N_0(t,v)t_\alpha v_2 t_\alpha^{-1} H_{\Delta,0}}} |N_0(t,v) \cap H_{\Delta,0} : N_0(t,v) \cap t_\alpha H_{\Delta,0} t_\alpha^{-1}| \chi^w(t_\alpha) f(vv_2)
\end{aligned}$$

since the double coset $(N_0(t,v) \cap H_{\Delta,0})1t_\alpha H_{\Delta,0}t_\alpha^{-1}$ contains $|N_0(t,v) \cap H_{\Delta,0} : N_0(t,v) \cap t_\alpha H_{\Delta,0} t_\alpha^{-1}|$ left cosets in $H_{\Delta,0}/t_\alpha H_{\Delta,0}t_\alpha^{-1}$. Assume first $N_0(t,v) \cap H_{\Delta,0} \not\subseteq t_\alpha H_{\Delta,0} t_\alpha^{-1}$ whence this number is divisible by p . In this case $F_\alpha(f) = 0$ for all $f \in \mathcal{C}_{w,0}(\chi)^{H_{\Delta,0}}$ since $p = 0$ in κ . So we deduce that F_α acts by 0. On the other hand, if $N_0(t,v) \cap H_{\Delta,0} \subseteq t_\alpha H_{\Delta,0} t_\alpha^{-1}$ and $\tilde{f}: N_{\Delta,0} \rightarrow \kappa$ denotes the function corresponding to f under the identification (10) then we compute

$$\widetilde{F_\alpha(f)}(\overline{v_1}) = \sum_{\substack{v_2 \in N_0(t,v) \setminus N_0/H_{\Delta,0} \\ v_1 \in N_0(t,v)t_\alpha v_2 t_\alpha^{-1} H_{\Delta,0}}} \chi^w(t_\alpha) f(vv_2) = \sum_{u \in J((N_{\Delta,0} \cap t_\alpha^{-1} N_{\Delta_1, *}) / N_{\Delta_1, *})} \chi(t_\alpha) \tilde{f}(t_\alpha^{-1} \overline{v_1} t_\alpha u)$$

for all $\overline{v_1} := v_1 H_{\Delta,0} \in N_{\Delta,0}$.

We denote by $\Delta' \subseteq \Delta$ the subset of those α simple roots satisfying $N_{t,v} > N_{t,t_\alpha v t_\alpha^{-1}}$ or $N_0(t,v) \cap H_{\Delta,0} \not\subseteq t_\alpha H_{\Delta,0} t_\alpha^{-1}$ so that F_α acts by 0 on $\text{gr}^{t,v}(\mathcal{C}_w)^{H_{\Delta,0}}$. Then the assumptions of Lemma 3.12 are satisfied and the statement follows. \square

Remark. Let $\chi: T \rightarrow \kappa^\times$ be a character and $w \in W$ such that for all roots $\beta \in (\Phi^+ \setminus \Delta) \cap w^{-1}(\Phi^-)$ and $\alpha \in \Delta \setminus w^{-1}(\Phi^-)$ we have $\beta \circ \alpha^\vee = 1$. Then the $\kappa[[N_{\Delta,0}]] [F_\Delta]$ -module $\mathcal{C}_{w,0}(\chi)^{H_{\Delta,0}}$ is finitely presented and simple. Otherwise the module $\mathcal{C}_{w,0}(\chi)^{H_{\Delta,0}}$ is not even finitely generated over $\kappa[[N_{\Delta,0}]] [F_\Delta]$.

Proposition 3.16. *Let $\chi: T \rightarrow \kappa^\times$ be a character. Then any finitely generated $\kappa[[N_{\Delta,0}]] [F_\Delta]$ -submodule M of $\mathcal{C}_w(\chi)^{H_{\Delta,0}}$ is finitely presented, has finite length, and is admissible as a representation of $N_{\Delta,0}$ for all $w \in W$.*

Proof. Any finitely generated $\kappa[[N_{\Delta,0}]] [F_\Delta]$ -submodule M of $\mathcal{C}_w(\chi)^{H_{\Delta,0}}$ is contained in $\text{Fil}^{t,v}$ for some $t \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ and $v \in N_0$. We prove the statement by induction on the number d of elements in $D_{t,w}$ that are smaller than $N_{t,v}$. If $d = 0$ then we have $v = 1$ and $\text{Fil}^{t,1} = \text{gr}^{t,1}(\mathcal{C}_w) = \mathcal{C}_{w,0}(\chi)$ whence the statement follows from Lemma 3.15. So assume $d > 0$. The image \overline{M} of M in $\text{gr}^{t,v}(\mathcal{C}_w) = \text{Fil}^{t,v} / \text{Fil}^{t,v'}$ is finitely presented and has finite length by Lemma 3.15 (here $N_{t,v'}$ is the biggest element in $D_{t,w}$ that is strictly smaller than $N_{t,v}$). Hence by Lemma 3.10(ii) $M \cap \text{Fil}^{t,v'}$ is finitely generated and the statement follows by induction on d using Lemma 3.10(i).

The admissibility of M follows from the fact that Fil^t is admissible as a representation of N_0 for each $t \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$. Indeed, $(\text{Fil}^t)^\vee$ is generated by at most $|t^{-1}N_0t : N_0|$ elements as a module over $\kappa[[N_0]]$. \square

Proposition 3.17. *Let π be an object in SP_A . Then any finitely generated $A[[N_{\Delta,0}]] [F_\Delta]$ -submodule M of $\pi^{H_{\Delta,0}}$ is finitely presented, has finite length, and is admissible as a representation of $N_{\Delta,0}$.*

Proof. This follows from Prop. 3.16 and Lemma 3.10 noting that each irreducible subquotient of $\pi|_B$ is isomorphic to $\mathcal{C}_w(\chi)$ for some character $\chi: T \rightarrow \kappa^\times$ and $w \in W$ by [27]. \square

Proposition 3.18. *Let π be an object in SP_A . The generic length of $D_\Delta^\vee(\pi)$ as a module over $A((N_{\Delta,0}))$ equals the number of Jordan-Hölder factors of $\pi|_B$ isomorphic to $\mathcal{C}_1(\chi)$ for some character $\chi: T \rightarrow \kappa^\times$. In particular, $D_\Delta^\vee(\pi)$ is finitely generated over $A((N_{\Delta,0}))$.*

Proof. We prove by induction on the length of $\pi|_B$ (that is finite by [27]). If $\pi|_B$ is irreducible then the statement follows combining Thm. 2.10 and Lemma 3.4 with Cor. 3.9. By [27] we have a short exact sequence $0 \rightarrow \pi_1 \rightarrow \pi \rightarrow \mathcal{C}_w(\chi) \rightarrow 0$ of B -representations for some character $\chi: T \rightarrow \kappa^\times$ and $w \in W$. By Thm. 2.10 this induces an exact sequence $D_\Delta^\vee(\mathcal{C}_w(\chi)) \rightarrow D_\Delta^\vee(\pi) \rightarrow D_\Delta^\vee(\pi_1) \rightarrow 0$. If $w \neq 1$ then we have $D_\Delta^\vee(\mathcal{C}_w(\chi)) = 0$ by Lemma 3.4 therefore there is nothing to prove. So assume $w = 1$. Then $D_\Delta^\vee(\mathcal{C}_1(\chi))$ has generic length 1 by Cor. 3.9. Moreover, $M := \mathcal{C}_{1,0}(\chi)^{H_{\Delta,0}}$ is finitely presented and simple as a module over $A[[N_{\Delta,0}]] [F_\Delta]$ by Lemma 3.15 and the Remark thereafter (as it is generated by the single $\mathbb{1}_{N_0}$) such that we have $M^\vee[1/X_\Delta] = D_\Delta^\vee(\mathcal{C}_1(\chi))$.

The $A[[X]] [F]$ -submodule M_ℓ of $\mathcal{C}_1(\chi)^{H_{\ell,0}}$ (see section 2.4 for the definition of $H_{\ell,0}$) generated by $\mathbb{1}_{N_0}$ is the unique minimal element in $\mathcal{M}(\mathcal{C}_1(\chi)^{H_{\ell,0}})$ with $M_\ell^\vee[1/X] \neq 0$. (To see this one could either use the same argument as in the proof of Lemma 3.12 to show that M_ℓ is simple or note that $D_\xi^\vee(\mathcal{C}_1(\chi)^{H_{\ell,0}})M_\ell^\vee[1/X]$ has dimension 1 over $\kappa((X))$ and M_ℓ^\vee is free of rank 1 over $\kappa[[X]]$ so we have $M_\ell^\vee[1/X] = 0$ for any proper submodule $M'_\ell \subsetneq M_\ell$. On the other hand, D_ξ^\vee is exact on the category SP_A (Cor. 9.2 in [8]). Therefore the natural map $M_\ell^\vee[1/X] \rightarrow D_\xi^\vee(\pi)$ is injective. So there exists an element $M_{1,\ell} \in \mathcal{M}(\pi^{H_{\ell,0}})$ whose image in $\mathcal{C}_1(\chi)^{H_{\ell,0}}$ contains M_ℓ . In particular, $\mathbb{1}_{N_0}$ is the image of an element $m \in \pi^{H_{\ell,0}} \subseteq \pi^{H_{\Delta,0}}$. Let us denote by M_1 the $A[[N_{\Delta,0}]] [F_\Delta]$ -submodule of $\pi^{H_{\Delta,0}}$ generated by the finite T_0 -orbit of m . By Prop. 3.17 M_1 is an object in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ admitting a surjective T_0 - and $A[[N_{\Delta,0}]] [F_\Delta]$ -equivariant map $M_1 \rightarrow M$. Therefore the composite map $M^\vee[1/X_\Delta] \cong D_\Delta^\vee(\mathcal{C}_1(\chi)) \rightarrow D_\Delta^\vee(\pi) \rightarrow M_1^\vee[1/X_\Delta]$ is injective so we have a short exact sequence $0 \rightarrow D_\Delta^\vee(\mathcal{C}_1(\chi)) \rightarrow D_\Delta^\vee(\pi) \rightarrow D_\Delta^\vee(\pi_1) \rightarrow 0$. \square

Theorem 3.19. *Let $0 \rightarrow \pi_1 \rightarrow \pi_2 \rightarrow \pi_3 \rightarrow 0$ be an exact sequence in SP_A . Then the sequence $0 \rightarrow D_\Delta^\vee(\pi_3) \rightarrow D_\Delta^\vee(\pi_2) \rightarrow D_\Delta^\vee(\pi_1) \rightarrow 0$ is also exact.*

Proof. By Thm. 2.10 we have an exact sequence $D_\Delta^\vee(\pi_3) \xrightarrow{f_3} D_\Delta^\vee(\pi_2) \rightarrow D_\Delta^\vee(\pi_1) \rightarrow 0$. By Prop. 3.18 we have $\text{length}_{\text{gen}}(D_\Delta^\vee(\pi_2)) = \text{length}_{\text{gen}}(D_\Delta^\vee(\pi_1)) + \text{length}_{\text{gen}}(D_\Delta^\vee(\pi_3))$. Hence we deduce $\text{length}_{\text{gen}}(\text{Im}(f_3)) = \text{length}_{\text{gen}}(D_\Delta^\vee(\pi_3))$ and $\text{length}_{\text{gen}}(\text{Ker}(f_3)) = 0$. By Prop. 2.5 $\text{Ker}(f_3) \cap D_\Delta^\vee(\pi_3)[\varpi]$ is a finitely generated and torsion étale T_+ -module over $\kappa((N_{\Delta,0}))$. Therefore its global annihilator is a T_0 -invariant ideal that equals $\kappa((N_{\Delta,0}))$ by Prop. 2.1. We deduce that f_3 is injective as desired. \square

Corollary 3.20. *Let π be an object in SP_A . Then we have $D_\xi^\vee(\pi) \cong A((X)) \otimes_{\ell, A((N_{\Delta,0}))} D_\Delta^\vee(\pi)$.*

Proof. We have a surjective map $D_\xi^\vee(\pi) \rightarrow A((X)) \otimes_{\ell, A((N_{\Delta,0}))} D_\Delta^\vee(\pi)$ by the universal property of D_ξ^\vee (see Remark 2 after Prop. 2.6) and the two sides have the same length as a module over the artinian ring $A((X))$ by Prop. 2.6 and Thm. 3.19. \square

4 Noncommutative theory

4.1 The ring $A((N_{\Delta,\infty}))$ and its first ring-theoretic properties

Let $H_{\Delta,k}$ be the normal subgroup of N_0 generated by $s^k H_{\Delta,0} s^{-k}$, ie. we put

$$H_{\Delta,k} = \langle n_0 s^k H_{\Delta,0} s^{-k} n_0^{-1} \mid n_0 \in N_0 \rangle .$$

$H_{\Delta,k}$ is an open subgroup of $H_{\Delta,0}$ normal in N_0 and we have $\bigcap_{k \geq 0} H_{\Delta,k} = \{1\}$. Put $N_{\Delta,k} := N_0/H_{\Delta,k}$ and consider the Iwasawa algebra $A[[N_{\Delta,k}]]$. There is a natural surjective homomorphism $N_{\Delta,k} \twoheadrightarrow N_{\Delta,0}$ with finite kernel $H_{\Delta,0}/H_{\Delta,k}$. This group homomorphism does not admit a splitting in general. However, there is a canonical splitting from the subgroup $s^k N_{\Delta,0} s^{-k} \leq N_{\Delta,0}$. Indeed, the image of the map $s^k(\cdot)s^{-k}: N_{\Delta,k} \rightarrow N_{\Delta,k}$ maps isomorphically onto $s^k N_{\Delta,0} s^{-k}$ since its intersection with the finite subgroup $H_{\Delta,0}/H_{\Delta,k}$ is trivial ($H_{\Delta,0}/H_{\Delta,k}$ is also the kernel of the homomorphism $s^k(\cdot)s^{-k}: N_{\Delta,k} \rightarrow N_{\Delta,k}$, so its image is torsion-free). Let $n_0 = n_0(G) \in \mathbb{N}$ be the maximum of the degrees of the algebraic characters $\beta \circ \xi: \mathbf{G}_m \rightarrow \mathbf{G}_m$ for all β in Φ^+ . Note that for any positive root $\beta \in \Phi^+$ we have $s N_{\beta,0} s^{-1} = N_{\beta,0}^{p^{\deg(\beta \circ \xi)}} \geq N_{\beta,0}^{p^{n_0}}$. Then $s^{kn_0} N_{\Delta,k} s^{-kn_0}$ even lies in the centre of the group $N_{\Delta,k}$: Since N_0 is totally decomposed, it suffices to verify that the commutator $[s^{kn_0} n_\alpha s^{-kn_0}, n_\beta]$ lies in $H_{\Delta,k}$ for the generators $n_\alpha := u_\alpha(1) \in N_{\alpha,0}$ and $n_\beta := u_\beta(1) \in N_{\beta,0}$ with $\alpha, \beta \in \Phi^+$. Now $s^{kn_0} n_\alpha s^{-kn_0}$ lies in $N_{\alpha,0}^{p^{kn_0 \alpha(s)}} \leq N_{\alpha,0}^{p^{kn_0}}$ therefore the commutator $[s^{kn_0} n_\alpha s^{-kn_0}, n_\beta]$ is in

$$[N_0, N_0]^{p^{kn_0}} \leq H_{\Delta,0}^{p^{kn_0}} = \langle N_{\beta,0}^{p^{kn_0}} \mid \beta \in \Phi^+ \setminus \Delta \rangle \leq \prod_{\beta \in \Phi^+ \setminus \Delta} s^k N_{\beta,0} s^{-k} = s^k H_{\Delta,0} s^{-k} \leq H_{\Delta,k} .$$

By an abuse of notation we also denote the class of n_α in $N_{\Delta,k}$ by n_α . Since $n_\alpha^{p^{kn_0}}$ lies in the centre of $N_{\Delta,k}$ we may form the ring

$$A((N_{\Delta,k})) := A[[N_{\Delta,k}]][(n_\alpha^{p^{kn_0}} - 1)^{-1} \mid \alpha \in \Delta] .$$

Note that $n_\alpha^{p^{kn_0}} - 1$ is divisible by $n_\alpha - 1$ therefore $n_\alpha - 1$ is also invertible in the ring $A((N_{\Delta,k}))$. Vice versa, if $n_\alpha - 1$ is invertible in a ring in which $p^h = 0$ then so is $n_\alpha^{p^{kn_0}} - 1$ since the latter differs from the invertible element $(n_\alpha - 1)^{p^{kn_0}}$ by something divisible by p which is nilpotent. So the ring $A((N_{\Delta,k}))$ does not depend on the choice of our power $n_\alpha^{p^{kn_0}}$, nor on the choice of the topological generator n_α . Moreover, for integers $0 \leq k_1 \leq k_2$ we have a natural surjective homomorphism

$$A((N_{\Delta,k_2})) \twoheadrightarrow A((N_{\Delta,k_1}))$$

induced by the group homomorphism $N_{\Delta,k_2} \twoheadrightarrow N_{\Delta,k_1}$.

Lemma 4.1. *For integers $0 \leq k_1 \leq k_2$ we have $A((N_{\Delta,k_1})) \cong H_0(H_{\Delta,k_1}/H_{\Delta,k_2}, A((N_{\Delta,k_2})))$ and $H_i(H_{\Delta,k_1}/H_{\Delta,k_2}, A((N_{\Delta,k_2}))) = 0$ for all $i > 0$. On finitely generated $A((N_{\Delta,k_2}))$ -modules the functors $H_0(H_{\Delta,k_1}/H_{\Delta,k_2}, \cdot)$ and $A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} \cdot$ are naturally isomorphic.*

Proof. For the completed group algebras we certainly have the analogous statements. More explicitly, since $H_{\Delta,k_1}/H_{\Delta,k_2}$ is a finite group, the kernel of the quotient map $A[[N_{\Delta,k_2}]] \rightarrow A[[N_{\Delta,k_1}]]$ is generated as a right (or left) ideal in $A[[N_{\Delta,k_2}]]$ by the elements $h-1$ ($h \in S$) for any fixed set S of generators of the group $H_{\Delta,k_1}/H_{\Delta,k_2} = \text{Ker}(N_{\Delta,k_2} \rightarrow N_{\Delta,k_1})$. Since localization is exact, the same is true for the kernel I of the natural map $A((N_{\Delta,k_2})) \rightarrow A((N_{\Delta,k_1}))$ showing that $A((N_{\Delta,k_1})) \cong H_0(H_{\Delta,k_1}/H_{\Delta,k_2}, A((N_{\Delta,k_2})))$. Now if M is a finitely generated module over $A((N_{\Delta,k_2}))$ then $H_0(H_{\Delta,k_1}/H_{\Delta,k_2}, M)$ is the quotient of M by the submodule generated by $(h-1)M$ ($h \in S$). We deduce

$$H_0(H_{\Delta,k_1}/H_{\Delta,k_2}, M) \cong M/IM \cong A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} M .$$

Finally, note that the completed group ring $A[[N_{\Delta,k_2}]]$ is a finite free module over its subring $A[[s^{k_2 n_0} N_{\Delta,k_2} s^{-k_2 n_0}]] [H_{\Delta,0}/H_{\Delta,k_2}]$. Inverting central elements $(n_\alpha^{p^{k_2 n_0}} - 1)$ for $\alpha \in \Delta$ we deduce that $A((N_{\Delta,k_2}))$ is finite free over $A((s^{k_2 n_0} N_{\Delta,k_2} s^{-k_2 n_0})) [H_{\Delta,0}/H_{\Delta,k_2}]$. In particular, it is induced as a representation of the finite group $H_{\Delta,0}/H_{\Delta,k_2}$ whence also as a representation of the subgroup $H_{\Delta,k_1}/H_{\Delta,k_2}$. Hence its higher homology groups vanish. \square

So we may form the projective limit

$$A((N_{\Delta,\infty})) := \varprojlim_k A((N_{\Delta,k})) .$$

Lemma 4.2. *The kernel of the natural map $A((N_{\Delta,\infty})) \rightarrow A((N_{\Delta,0}))$ is generated as a left (or as a right) ideal by the elements $n_\beta - 1$ for topological generators $n_\beta \in N_{\beta,0}$ for $\beta \in \Phi^+ \setminus \Delta$.*

Proof. We prove by induction on $\dim N_0$ as a p -adic Lie group. Note that N_0 is nilpotent so unless $\Delta = \Phi^+$ (whence there is nothing to prove), we have a β in $\Phi^+ \setminus \Delta$ such that $N_{\beta,0}$ lies in the centre of N_0 . Then we may form the ring $A((N_{\Delta,\infty,\beta}))$ with the group N_0 replaced by $N_0/N_{\beta,0}$. For the k th layer it is clear that the kernel of the natural map $A((N_{\Delta,k})) \rightarrow A((N_{\Delta,k,\beta}))$ is generated by $n_\beta - 1$. Moreover, the maps $A((N_{\Delta,k}))(n_\beta - 1) \rightarrow A((N_{\Delta,k-1}))(n_\beta - 1)$ are clearly surjective. Therefore the projective limit of the diagrams

$$0 \rightarrow A((N_{\Delta,k}))(n_\beta - 1) \rightarrow A((N_{\Delta,k})) \rightarrow A((N_{\Delta,k,\beta})) \rightarrow 0$$

is exact whence the kernel of the map $A((N_{\Delta,\infty})) \rightarrow A((N_{\Delta,\infty,\beta}))$ is also generated by $n_\beta - 1$. Now we have a composite map

$$A((N_{\Delta,\infty})) \xrightarrow{\Pi_\beta} A((N_{\Delta,\infty,\beta})) \xrightarrow{\Pi_{\Phi^+ \setminus (\Delta \cup \{\beta\})}} A((N_{\Delta,0})) .$$

We have just shown that $\text{Ker}(\Pi_\beta)$ is generated by $n_\beta - 1$ and by the inductive hypothesis we see that $\text{Ker}(\Pi_{\Phi^+ \setminus (\Delta \cup \{\beta\})})$ is generated by $\{n_\gamma - 1 \mid \gamma \in \Phi^+ \setminus (\Delta \cup \{\beta\})\}$. The statement follows by combining these two. \square

We denote by φ_t the ring endomorphism of $A((N_{\Delta,k}))$ induced by the conjugation of $t \in T_+$ on $N_{\Delta,k}$ for all $k \geq 0$. This induces a ring endomorphism (still denoted by φ_t) on $A((N_{\Delta,\infty}))$ by taking the projective limit.

Lemma 4.3. For each $t \in T_+$ the ring homomorphism $\varphi_t: A((N_{\Delta, \infty})) \rightarrow A((N_{\Delta, \infty}))$ is injective and we have

$$A((N_{\Delta, \infty})) = \bigoplus_{u \in J(N_0/tN_0t^{-1})} u\varphi_t(A((N_{\Delta, \infty}))) = \bigoplus_{u \in J(N_0/tN_0t^{-1})} \varphi_t(A((N_{\Delta, \infty})))u$$

as left (resp. right) modules over the the subring $\varphi_t(A((N_{\Delta, \infty})))$.

Proof. Let $t \in T_+$ be arbitrary and choose an integer $k(t) > 0$ so that we have $H_{\Delta, k(t)} \leq tN_0t^{-1}$ and $s^{k(t)n_0}t^{-1} \in T_+$, and let $k \geq k(t)$. Then we have $|N_0 : \varphi_t(N_0)| = |N_{\Delta, k} : \varphi_t(N_{\Delta, k})|$. In particular, for the Iwasawa algebra $A[[N_{\Delta, k}]]$ we have

$$A[[N_{\Delta, k}]] = \bigoplus_{u \in J(N_0/tN_0t^{-1})} u\varphi_t(A[[N_{\Delta, k}]]) . \quad (12)$$

By the choice of k the element $s^{kn_0}n_\alpha s^{-kn_0} - 1$ lies in the subring $\varphi_t(A[[N_{\Delta, k}]])$ for each $\alpha \in \Delta$. So inverting $s^{kn_0}n_\alpha s^{-kn_0} - 1$ for each $\alpha \in \Delta$ on both sides of (12) we obtain

$$\begin{aligned} A((N_{\Delta, k})) &= \bigoplus_{u \in J(N_0/tN_0t^{-1})} u\varphi_t(A[[N_{\Delta, k}]][\varphi_{s^{kn_0}t^{-1}}(n_\alpha - 1)^{-1} \mid \alpha \in \Delta]) = \\ &= \bigoplus_{u \in J(N_0/tN_0t^{-1})} u\varphi_t(A((N_{\Delta, k}))) \end{aligned} \quad (13)$$

since by inverting $\varphi_{s^{kn_0}t^{-1}}(n_\alpha - 1)$ the element $\varphi_{s^{kn_0}t^{-1}}(n_\alpha - 1)$ becomes invertible, too, and vice versa. We deduce the statement by taking projective limits. The left module version follows similarly.

For the injectivity of φ_t on $A((N_{\Delta, \infty}))$ note that we have a short exact sequence

$$0 \rightarrow (t^{-1}H_{\Delta, k}t - 1)A[[N_{\Delta, k}]] \rightarrow A[[N_{\Delta, k}]] \xrightarrow{\varphi_t} \varphi_t(A[[N_{\Delta, k}]]) \rightarrow 0 .$$

Inverting again the central elements $\varphi_{s^{kn_0}t^{-1}}(n_\alpha - 1)$ for each $\alpha \in \Delta$ we obtain a projective system of short exact sequences

$$0 \rightarrow (t^{-1}H_{\Delta, k}t - 1)A((N_{\Delta, k})) \rightarrow A((N_{\Delta, k})) \xrightarrow{\varphi_t} \varphi_t(A((N_{\Delta, k}))) \rightarrow 0 . \quad (14)$$

Now there exists an integer $k_2 > k$ such that $H_{\Delta, k_2} \leq tH_{\Delta, k}t^{-1}$ whence $(t^{-1}H_{\Delta, k_2}t - 1) \subseteq H_{\Delta, k} - 1$. So the connecting map for the left hand side of (14) is the zero map from the k_2 th level to the k th level. By taking projective limits we obtain the required injectivity. \square

Proposition 4.4. The ring $A((N_{\Delta, \infty}))$ is (left and right) noetherian.

Proof. We may assume without loss of generality that N_0 is uniform as a pro- p group since any ring that is finitely generated as a module over a noetherian subring is itself noetherian. Moreover, it suffices to treat the case $h = 1$. We consider the filtration on $\kappa((N_{\Delta, \infty}))$ by the powers of the ideal $I := \text{Ker}(\kappa((N_{\Delta, \infty})) \rightarrow \kappa((N_{\Delta, 0})))$. This filtration is cofinal with the filtration induced by the kernels of the maps $\kappa((N_{\Delta, \infty})) \rightarrow \kappa((N_{\Delta, n}))$ ($n \geq 0$) therefore $\kappa((N_{\Delta, \infty}))$ is complete with respect to the I -adic topology. By Lemma 4.2 the powers I^n of the ideal I are generated by n -term products of the elements $n_\beta - 1$ for $\beta \in \Phi^+ \setminus \Delta$. Since N_0

is uniform, its commutator subgroup is contained in N_0^p . Moreover, the commutator $[N_0, N_0]$ of N_0 is also contained in $H_{\Delta,0}$ since $N_0/H_{\Delta,0} \cong N_{\Delta,0}$ is commutative. Since $N_{\Delta,0}$ does not have elements of order p , $[N_0, N_0]$ is contained in $H_{\Delta,0}^p$. This shows that the graded ring $\text{gr} \kappa((N_{\Delta,\infty}))$ of $\kappa((N_{\Delta,\infty}))$ is a commutative polynomial ring over $\kappa((N_{\Delta,0}))$ in the variables $X_\beta = \text{gr}(n_\beta - 1)$ for $\beta \in \Phi^+ \setminus \Delta$. This is a noetherian ring by Hilbert's basis theorem as $\kappa((N_{\Delta,0}))$ is the localization of the noetherian ring $\kappa[[N_{\Delta,0}]]$ therefore also noetherian. Finally, the statement follows from Prop. I.7.1.2 in [22]. \square

Lemma 4.5. *The Jacobson radical of $A((N_{\Delta,\infty}))$ is the ideal generated by ϖ and $(H_{\Delta,0} - 1)A((N_{\Delta,\infty}))$.*

Proof. $A((N_{\Delta,\infty}))$ is complete with respect to the filtration induced by the powers of $\varpi A((N_{\Delta,\infty})) + (H_{\Delta,0} - 1)A((N_{\Delta,\infty}))$. Therefore this ideal is contained in the Jacobson radical of $A((N_{\Delta,\infty}))$. The other direction follows from noting that the Jacobson radical of $\kappa((N_{\Delta,0})) \cong \kappa[[X_\alpha \mid \alpha \in \Delta]][X_\alpha^{-1} \mid \alpha \in \Delta]$ is 0 by Prop. 2.1 since it is T_0 -invariant. \square

4.2 The equivalence of categories

By Lemma 4.3 we may consider étale T_+ -modules over $A((N_{\Delta,\infty}))$ the usual way: an étale T_+ -module D_∞ over $A((N_{\Delta,\infty}))$ is a finitely generated $A((N_{\Delta,\infty}))$ -module with a semilinear action of the monoid T_+ such that for all $t \in T_+$ the map

$$\begin{aligned} 1 \otimes \varphi_t: A((N_{\Delta,\infty})) \otimes_{\varphi_t, A((N_{\Delta,\infty}))} D_\infty &\rightarrow D_\infty \\ \lambda \otimes x &\mapsto \lambda \varphi_t(x) \end{aligned}$$

is an isomorphism. Similarly, we may consider étale T_* -modules over $A((N_{\Delta,\infty}))$ for any submonoid $T_* \leq T_+$ and also over the rings $A((N_{\Delta,k}))$ for each $k \geq 0$.

Let $T_* \leq T_+$ be a submonoid containing some power s^{r_*} of $s = \xi(p)$. Our goal is to prove an equivalence between the category $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ of (finitely generated) étale T_* -modules over $A((N_{\Delta,0}))$ and the category $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$ of (finitely generated) étale T_* -modules over $A((N_{\Delta,\infty}))$. The reason why this is not a formal consequence of Thm. 8.20 in [26] or of Prop. 3.1 in [28] or even of those proofs is that there is no section of the group homomorphism $N_0 \twoheadrightarrow N_{\Delta,0}$ in general. Therefore it is not obvious a priori how to construct a functor from $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ to $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$. The idea is to prove an equivalence over each level k and to take the projective limit. Even though there is no section of the group homomorphism $N_{\Delta,k} \twoheadrightarrow N_{\Delta,0}$ either, we do have a section from a finite index subgroup $s^k N_{\Delta,0} s^{-k}$.

For an étale T_* -module D over $A((N_{\Delta,0}))$ and integer $r_* \mid k$ note that we have a ring homomorphism $\varphi_{s^{kn_0}}: A((N_{\Delta,0})) \hookrightarrow A((s^{kn_0} N_{\Delta,0} s^{-kn_0})) \leq A((N_{\Delta,k}))$. So we define

$$D_k := \mathbb{M}_{k,0}(D) := A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D .$$

This is an étale T_* -module over $A((N_{\Delta,k}))$ as we compute

$$\begin{aligned} A((N_{\Delta,k})) \otimes_{\varphi_t, A((N_{\Delta,k}))} D_k &= A((N_{\Delta,k})) \otimes_{\varphi_t, A((N_{\Delta,k}))} A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D \cong \\ &\cong A((N_{\Delta,k})) \otimes_{\varphi_{t s^{kn_0}}, A((N_{\Delta,0}))} D \cong A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} A((N_{\Delta,0})) \otimes_{\varphi_t, A((N_{\Delta,0}))} D \cong \\ &\cong A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D \cong D_k \end{aligned}$$

for each $t \in T_*$. The above identification is given by the map $1 \otimes \varphi_t : A((N_{\Delta,k})) \otimes_{\varphi_t, A((N_{\Delta,k}))} D_k \rightarrow D_k$. On the other hand, we compute

$$\begin{aligned} H_0(H_{\Delta,0}/H_{\Delta,k}, D_k) &= H_0(H_{\Delta,0}/H_{\Delta,k}, A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D) \cong \\ &\cong H_0(H_{\Delta,0}/H_{\Delta,k}, A((N_{\Delta,k}))) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D \cong A((N_{\Delta,0})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} D \cong D \end{aligned}$$

by Lemma 4.1. So we obtained a natural isomorphism between the identity functor on $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ and the functor $H_0(H_{\Delta,0}/H_{\Delta,k}, (\cdot)_k)$.

Lemma 4.6. *The functors $\mathbb{M}_{k,0} : D \mapsto D_k$ and $\mathbb{D}_{0,k} : D_k \mapsto H_0(H_{\Delta,0}/H_{\Delta,k}, D_k)$ are quasi-inverse equivalences of categories between $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ and $\mathcal{D}^{et}(T_*, A((N_{\Delta,k})))$.*

Proof. We have already seen that $\mathbb{D}_{0,k} \circ \mathbb{M}_{k,0} \cong \text{id}$. For the other direction let D_k be an object in $\mathcal{D}^{et}(T_*, A((N_{\Delta,k})))$. Note that by Lemma 4.1 we have $\mathbb{D}_{0,k} = A((N_{\Delta,0})) \otimes_{A((N_{\Delta,k}))} \cdot$. Moreover, $(H_{\Delta,0}/H_{\Delta,k} - 1)A((N_{\Delta,k}))$ lies in the kernel of the ring homomorphism $\varphi_{s^{kn_0}} : A((N_{\Delta,k})) \rightarrow A((N_{\Delta,0}))$. So we may factor $\varphi_{s^{kn_0}}$ as

$$\begin{array}{ccc} A((N_{\Delta,k})) & \longrightarrow & A((N_{\Delta,0})) \xrightarrow{\varphi_{s^{kn_0}}} A((N_{\Delta,k})) \\ & \searrow \text{curved arrow} & \nearrow \\ & & \varphi_{s^{kn_0}} \end{array}$$

Therefore we compute

$$\begin{aligned} \mathbb{M}_{k,0} \circ \mathbb{D}_{0,k}(D_k) &= A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,0}))} A((N_{\Delta,0})) \otimes_{A((N_{\Delta,k}))} D_k \cong \\ &\cong A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}}, A((N_{\Delta,k}))} D_k \cong D_k \end{aligned}$$

where the last isomorphism follows from the étaleness of D_k . \square

For an object D in $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ and integers $k_1 \leq k_2$ both divisible by r_* we have

$$\mathbb{M}_{k_1,0}(D) = A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} \mathbb{M}_{k_2,0}(D) .$$

In particular, $(\mathbb{M}_{k,0}(D))_{k \geq 0}$ forms a projective system. So we define $\mathbb{M}_{\infty,0} := \varprojlim_k \mathbb{M}_{k,0}$ and $\mathbb{D}_{0,\infty} := H_0(H_{\Delta,0}, \cdot) = A((N_{\Delta,0})) \otimes_{A((N_{\Delta,\infty}))} \cdot$. $\mathbb{D}_{0,\infty}$ is a functor from $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$ to $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$. It is not so trivial (but as we shall see, true) that $\mathbb{M}_{\infty,0}$ is also a functor in the reverse direction. Note that any module over $A((N_{\Delta,k}))$ can be regarded as a module over $A((N_{\Delta,\infty}))$ via the quotient map $A((N_{\Delta,\infty})) \twoheadrightarrow A((N_{\Delta,k}))$ for any $k \geq 0$ divisible by r_* . Moreover, we have a semilinear action of T_* on each $\mathbb{M}_{k,0}(D)$ whence also on the projective limit as the connecting maps are T_* -equivariant by construction. In particular, $\mathbb{M}_{\infty,0}(D)$ is a module over $A((N_{\Delta,\infty}))$ with a semilinear action of T_* . Our key result is the following

Proposition 4.7. *For each object D in $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ the $A((N_{\Delta,\infty}))$ -module $\mathbb{M}_{\infty,0}(D)$ with the T_* -action induced by the T_* -action on D is an object in $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$.*

Proof. We proceed in 3 steps.

Step 1: We show that $\mathbb{M}_{\infty,0}(D)$ is finitely generated over $A((N_{\Delta,\infty}))$. Note that the kernel of the ring homomorphism $A((N_{\Delta,k})) \twoheadrightarrow A((N_{\Delta,0}))$ is a nilpotent ideal therefore contained in the Jacobson radical of $A((N_{\Delta,k}))$. So if D is generated by the elements d_1, \dots, d_r then any

lifts $d_{1,k}, \dots, d_{r,k}$ of d_1, \dots, d_r to $\mathbb{M}_{k,0}(D)$ generate $\mathbb{M}_{k,0}(D)$ by Nakayama's Lemma ($r_* \mid k$). Since the natural quotient maps $\mathbb{M}_{k_2,0}(D) \rightarrow \mathbb{M}_{k_1,0}(D)$ are surjective for each pair $k_1 \leq k_2$ of integers (both divisible by r_*), we can choose the lifts $d_{1,k}, \dots, d_{r,k}$ recursively in a compatible way so that d_{i,k_2} maps to d_{i,k_1} under the quotient map $\mathbb{M}_{k_2,0}(D) \rightarrow \mathbb{M}_{k_1,0}(D)$. Now for $k_1 \leq k_2$ consider the commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{k_2} & \longrightarrow & \bigoplus_{i=1}^r A((N_{\Delta,k_2}))e_{i,k_2} & \xrightarrow{e_{i,k_2} \mapsto d_{i,k_2}} & \mathbb{M}_{k_2,0}(D) \longrightarrow 0 \\
& & \downarrow & & \downarrow e_{i,k_2} \mapsto e_{i,k_1} & & \downarrow \\
0 & \longrightarrow & C_{k_1} & \longrightarrow & \bigoplus_{i=1}^r A((N_{\Delta,k_1}))e_{i,k_1} & \xrightarrow{e_{i,k_1} \mapsto d_{i,k_1}} & \mathbb{M}_{k_1,0}(D) \longrightarrow 0
\end{array} \quad (15)$$

in which C_{k_j} ($j = 1, 2$) are defined so as to make the rows exact. Since the map $\mathbb{M}_{k_2,0}(D) \rightarrow \mathbb{M}_{k_1,0}(D)$ is induced by the identification $\mathbb{M}_{k_1,0}(D) \cong A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} \mathbb{M}_{k_2,0}(D)$ and the horizontal map in the middle is also induced by the natural quotient map $A((N_{\Delta,k_2})) \rightarrow A((N_{\Delta,k_1}))$, we obtain an exact sequence

$$A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} C_{k_2} \longrightarrow \bigoplus_{i=1}^r A((N_{\Delta,k_1}))e_{i,k_1} \xrightarrow{e_{i,k_1} \mapsto d_{i,k_1}} \mathbb{M}_{k_1,0}(D) \longrightarrow 0. \quad (16)$$

We deduce that the composite map $C_{k_2} \twoheadrightarrow A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} C_{k_2} \rightarrow C_{k_1}$ is also surjective. Therefore by the Mittag-Leffler condition we obtain an exact sequence

$$0 \longrightarrow \varprojlim_{r_* \mid k} C_k \longrightarrow \bigoplus_{i=1}^r A((N_{\Delta,\infty}))e_{i,\infty} \longrightarrow \mathbb{M}_{\infty,0}(D) \longrightarrow 0.$$

In particular, $\mathbb{M}_{\infty,0}(D)$ is finitely generated over $A((N_{\Delta,\infty}))$.

Step 2: We show that the map (18) below is injective for all $t \in T_$.*

Lemma 4.8. *For all $k \geq 0$ divisible by r_* We have $H_i(H, \mathbb{M}_{k,0}(D)) = 0$ for all $i > 0$ and subgroup $H \leq H_{\Delta,0}/H_{\Delta,k}$.*

Proof. The group ring $A((s^{kn_0} N_{\Delta,0} s^{-kn_0}))[H_{\Delta,0}/H_{\Delta,k}]$ is a subring in $A((N_{\Delta,k}))$. Moreover, we have

$$A((N_{\Delta,k})) = \bigoplus_{u \in J(N_0/s^{kn_0} N_0 s^{-kn_0} H_{\Delta,0})} A((s^{kn_0} N_{\Delta,0} s^{-kn_0}))[H_{\Delta,0}/H_{\Delta,k}]u.$$

In particular, $A((N_{\Delta,k}))$ is a free left module over $A((s^{kn_0} N_{\Delta,0} s^{-kn_0}))[H_{\Delta,0}/H_{\Delta,k}]$. Therefore $\mathbb{M}_{k,0}(D) = A((N_{\Delta,k})) \otimes_{\varphi_{s^{kn_0}, A((N_{\Delta,0}))}} D$ is an induced module as a representation of any subgroup $H \leq H_{\Delta,0}/H_{\Delta,k}$. In particular, its higher homology groups vanish. \square

Now we take $H_{\Delta,k_1}/H_{\Delta,k_2}$ -homology of the upper row in (15) and apply Lemma 4.8 (with $H := H_{\Delta,k_1}/H_{\Delta,k_2} \leq H_{\Delta,0}/H_{\Delta,k_2}$) to deduce the exactness of the sequence (16) on the left using Lemma 4.1 and the long exact sequence of homology. By the lower row in (15) we obtain the isomorphism $C_{k_1} \cong A((N_{\Delta,k_1})) \otimes_{A((N_{\Delta,k_2}))} C_{k_2}$ for each pair $k_1 \leq k_2$ (divisible by r_*).

Now for any fixed $k \geq 0$ we get a commutative diagram

$$\begin{array}{ccc}
& & 0 \\
& & \downarrow \\
A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} \varprojlim_{k'} C_{k'} & \longrightarrow & C_k \\
\downarrow & & \downarrow \\
\bigoplus_{i=1}^r A((N_{\Delta,k})) e_{i,\infty} & \longrightarrow & \bigoplus_{i=1}^r A((N_{\Delta,k})) e_{i,k} \\
\downarrow & & \downarrow \\
A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} \mathbb{M}_{\infty,0}(D) & \longrightarrow & \mathbb{M}_{k,0}(D) \\
\downarrow & & \downarrow \\
0 & & 0
\end{array} \tag{17}$$

with exact columns. The horizontal maps are onto since for example C_k is quotient of $\varprojlim_{k'} C_{k'}$ by the surjectivity of the maps $C_{k'} \rightarrow C_k$ ($k \leq k'$) and this quotient map factors through the maximal quotient of $\varprojlim_{k'} C_{k'}$ on which $A((N_{\Delta,\infty}))$ acts via its quotient $A((N_{\Delta,k}))$. Therefore the lower horizontal map is an isomorphism as the middle map clearly is.

Now take an arbitrary t in T_* . Since T_* acts on $\mathbb{M}_{\infty,0}(D)$, we have a map

$$\begin{aligned}
1 \otimes \varphi_t: A((N_{\Delta,\infty})) \otimes_{\varphi_t, A((N_{\Delta,\infty}))} \mathbb{M}_{\infty,0}(D) &\rightarrow \mathbb{M}_{\infty,0}(D) \\
\sum_{u \in J(N_0/tN_0t^{-1})} u \otimes m_u &\mapsto \sum_{u \in J(N_0/tN_0t^{-1})} u \varphi_t(m_u).
\end{aligned} \tag{18}$$

Write each m_u ($u \in J(N_0/tN_0t^{-1})$) as a sequence $m_u = (m_{u,k})_k$ with $m_{u,k} \in \mathbb{M}_{k,0}(D)$. Assume that $\sum_{u \in J(N_0/tN_0t^{-1})} u \otimes m_u$ lies in the kernel of the map (18). Then for each $k \geq 0$ (divisible by r_*) we have

$$\sum_{u \in J(N_0/tN_0t^{-1})} u \varphi_t(m_{u,k}) = 0.$$

Since $\mathbb{M}_{k,0}(D)$ is an étale T_* -module by Lemma 4.6, we obtain that $\sum_{u \in J(N_0/tN_0t^{-1})} u \otimes m_{u,k} = 0$ in $A((N_{\Delta,k})) \otimes_{\varphi_t, A((N_{\Delta,k}))} \mathbb{M}_{k,0}(D)$. Let now k be big enough so that $H_{\Delta,k}$ is contained in tN_0t^{-1} . Then $A((N_{\Delta,k}))$ is a free right module over the image of φ_t with generators $u \in J(N_0/tN_0t^{-1})$. Therefore we have $1 \otimes m_{u,k} = 0$ for each $u \in J(N_0/tN_0t^{-1})$ which shows that $m_{u,k}$ lies in $(t^{-1}H_{\Delta,k}t - 1)\mathbb{M}_{k,0}(D)$. So for any $0 \leq k' \leq k$ for which $H_{\Delta,k'}$ contains $t^{-1}H_{\Delta,k}t$ we deduce that $m_{u,k'} = 0$. However, for any fixed $0 \leq k'$ there exists a large enough $k \geq k'$ with $H_{\Delta,k'} \supseteq t^{-1}H_{\Delta,k}t$ so we have $m_{u,k'} = 0$ for all $k' \geq 0$. Therefore (18) is injective.

Step 3: We show that (18) is surjective. Let $m = (m_k)_k \in \mathbb{M}_{\infty,0}(D)$ be arbitrary. Since $\mathbb{M}_{k,0}(D)$ is an étale T_* -module over $A((N_{\Delta,k}))$ for all $k \geq 0$ divisible by r_* (see Lemma 4.6), we may decompose m_k as

$$m_k = \sum_{u \in J(N_0/tN_0t^{-1})} u \varphi_t(m_{u,k}) \tag{19}$$

for some $m_{u,k} \in \mathbb{M}_{k,0}(D)$. Note that the elements $m_{u,k}$ are not unique since φ_t is not injective on $\mathbb{M}_{k,0}(D)$. However, their images $\varphi_t(m_{u,k})$ under φ_t are unique for each $u \in J(N_0/tN_0t^{-1})$

and $k \geq k(t)$ large enough so that $H_{\Delta,k}$ is contained in tN_0t^{-1} . Fix $m_{u,k}$ arbitrarily for each $u \in J(N_0/tN_0t^{-1})$ and $k \geq k(t)$ (divisible by r_*) so that they satisfy (19) and put

$$Y_{u,k} := \{m'_{u,k} \in \mathbb{M}_{k,0}(D) \mid \varphi_t(m'_{u,k}) = \varphi_t(m_{u,k})\} .$$

Let $k \geq k(t)$ be fixed now let $k' \geq k$ be another integer so that we have $t^{-1}H_{\Delta,k'}t \leq H_{\Delta,k}$. The kernel of $\varphi_t: \mathbb{M}_{k',0}(D) \rightarrow \mathbb{M}_{k,0}(D)$ equals $(t^{-1}H_{\Delta,k'}t - 1)\mathbb{M}_{k',0}$. However, by our assumption that $t^{-1}H_{\Delta,k'}t$ is contained in $H_{\Delta,k}$, it follows that $(t^{-1}H_{\Delta,k'}t - 1)\mathbb{M}_{k',0}$ maps to 0 in $\mathbb{M}_{k,0}(D)$. Hence for each $u \in J(N_0/tN_0t^{-1})$ the image of $Y_{u,k}$ in $Y_{u,k}$ is a single element $m_{u,k}^* \in Y_{u,k}$. Therefore the projective system $(Y_{u,k})_k$ satisfies the Mittag-Leffler condition so that $Y_{u,\infty} := \varprojlim_k Y_{u,k} \subset \mathbb{M}_{\infty,0}(D)$ is a set having a single element $m_u := (m_{u,k}^*)_{k \geq 0}$. We clearly have $m = \sum_{u \in J(N_0/tN_0t^{-1})} u\varphi_t(m_u)$ as required. \square

Now our main result in this section becomes a simple application of the above Proposition.

Theorem 4.9. *The functors $\mathbb{M}_{\infty,0}$ and $\mathbb{D}_{0,\infty}$ are quasi-inverse equivalences of categories between $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$ and $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$.*

Proof. We saw in the proof of Prop. 4.7 that the lower horizontal map in (17) is an isomorphism for all $k \geq 0$ divisible by r_* . In the special case of $k = 0$ this provides us with a natural isomorphism between the identity and $\mathbb{D}_{0,\infty} \circ \mathbb{M}_{\infty,0}$ on $\mathcal{D}^{et}(T_*, A((N_{\Delta,0})))$.

Let D_∞ be an object in $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$. For each $k \geq 0$ (divisible by r_*) we have a natural quotient map

$$D_\infty \rightarrow D_k := A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} D_\infty .$$

Moreover, D_k is an étale T_* -module over $A((N_{\Delta,k}))$ corresponding to $D_0 = \mathbb{D}_{0,\infty}(D_\infty)$ via the equivalence of categories in Lemma 4.6. These reduction maps are compatible therefore we obtain a natural map $f: D_\infty \rightarrow \varprojlim_k D_k = \mathbb{M}_{\infty,0} \circ \mathbb{D}_{0,\infty}(D_\infty)$. This map f is an isomorphism modulo the Jacobson radical of $A((N_{\Delta,\infty}))$ by Lemma 4.5 using that $\mathbb{D}_{0,\infty} \circ \mathbb{M}_{\infty,0} \circ \mathbb{D}_{0,\infty} \cong \mathbb{D}_{0,\infty}$. In particular f is surjective. Moreover, $A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} f$ is also an isomorphism for all $k \geq 0$ divisible by r_* . Therefore we have $\text{Ker}(f) \subseteq (H_{\Delta,k} - 1)D_\infty$ for all k . Since the powers of the Jacobson radical J of $A((N_{\Delta,\infty}))$ are cofinal with the ideals $(H_{\Delta,k} - 1)A((N_{\Delta,\infty}))$ we deduce that $\text{Ker}(f) \subseteq J^k D_\infty$ for all $k \geq 0$. Now note that J is generated by a centralizing sequence by Lemma 4.2. Therefore it satisfies the Artin-Rees property by Thm. 4.2.7 in [23] and Prop. 4.4. This shows that $\bigcap_{k \geq 0} J^k D_\infty = \{0\}$ since D_∞ is finitely generated. In particular, f is an isomorphism. \square

Remark. The above proof shows that we have $D_\infty \cong \varprojlim_k A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} D_\infty$ for any object D_∞ in $\mathcal{D}^{et}(T_*, A((N_{\Delta,\infty})))$.

4.3 A noncommutative variant of D_Δ^\vee

The goal in this section is to define a noncommutative variant $D_{\Delta,\infty}^\vee$ of the functor D_Δ^\vee from smooth representations π of G over A to (projective limits of) étale T_+ -modules over $A((N_{\Delta,\infty}))$. The motivation is that this allows us to construct a G -equivariant sheaf on G/B (in the sense of [26]) attached to π which is crucial in possibly reconstructing π from $D_\Delta^\vee(\pi)$ in the case of extensions of principal series.

Let π be a smooth representation of B_0 over A together with an action of the monoid B_+ on π extending the action of B_0 . (For instance, π could be a smooth representation of B .) Denote by $F_{\alpha,k}$ the operator $\text{Tr}_{H_{\Delta,k}/t_\alpha H_{\Delta,k} t_\alpha^{-1}} \circ (t_\alpha \cdot)$ on $\pi^{H_{\Delta,k}}$ and consider the skew polynomial ring

$$A[[N_{\Delta,k}]] [F_{\Delta,k}] := A[[N_{\Delta,k}]] [F_{\alpha,k} \mid \alpha \in \Delta]$$

in the variables $F_{\alpha,k}$ ($\alpha \in \Delta$) that commute with each other and satisfy $F_{\alpha,k} \lambda = \varphi_\alpha(\lambda) F_{\alpha,k}$ for any $\lambda \in A[[N_{\Delta,k}]]$ and $\alpha \in \Delta$. Further, for any $t = \prod_{\alpha \in \Delta} t_\alpha^{k_\alpha}$ we put $F_{t,k} := \prod_{\alpha \in \Delta} F_{t_\alpha}^{k_\alpha}$. We denote by $\mathcal{M}_{\Delta,k}(\pi^{H_{\Delta,k}})$ the set of finitely generated $A[[N_{\Delta,k}]] [F_{\Delta,k}]$ -submodules of $\pi^{H_{\Delta,k}}$ that are stable under the action of T_0 and admissible as a representation of $N_{\Delta,k}$. We proceed as in section 2 of [17].

Lemma 4.10. *For each $\alpha \in \Delta$ we have $F_{\alpha,0} = F_\alpha$ and $F_{\alpha,k} \circ \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) = \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) \circ F_{\alpha,0}$ as maps on $\pi^{H_{\Delta,0}}$.*

Proof. We compute

$$\begin{aligned} & F_{\alpha,k} \circ \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/t_\alpha H_{\Delta,k} t_\alpha^{-1}} \circ (t_\alpha \cdot) \circ \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/t_\alpha H_{\Delta,k} t_\alpha^{-1}} \circ \text{Tr}_{t_\alpha H_{\Delta,k} t_\alpha^{-1}/s^k t_\alpha H_{\Delta,0} t_\alpha^{-1} s^{-k}} \circ (s^k t_\alpha \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/s^k t_\alpha H_{\Delta,0} t_\alpha^{-1} s^{-k}} \circ (s^k t_\alpha \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ \text{Tr}_{s^k H_{\Delta,0} s^{-k}/s^k t_\alpha H_{\Delta,0} t_\alpha^{-1} s^{-k}} \circ (s^k t_\alpha \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) \circ \text{Tr}_{H_{\Delta,0}/t_\alpha H_{\Delta,0} t_\alpha^{-1}} \circ (t_\alpha \cdot) = \\ &= \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k \cdot) \circ F_{\alpha,0} . \end{aligned}$$

□

Let M_0 be any (finitely generated) $A[[N_{\Delta,0}]] [F_\Delta]$ -module. Then $A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}], \varphi_{s^k}]} M_0$ naturally has the structure of a module over $A[[N_{\Delta,k}]] [F_{\Delta,k}]$ by putting $F_{\alpha,k}(\lambda \otimes m) := \varphi_{t_\alpha}(\lambda) \otimes F_\alpha(m)$. Let M be in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. In view of Lemma 4.10 we define M_k to be the image of the $A[[N_{\Delta,k}]] [F_{\Delta,k}]$ -module homomorphism

$$\begin{aligned} A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}], \varphi_{s^k}]} M &\rightarrow \pi^{H_{\Delta,k}} \\ \lambda \otimes m &\mapsto \lambda \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}}(s^k m) . \end{aligned}$$

In particular, M_k is an $A[[N_{\Delta,k}]] [F_{\Delta,k}]$ -submodule of $\pi^{H_{\Delta,k}}$. $\text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k M)$ is a $s^k N_0 s^{-k} H_{\Delta,k}$ -subrepresentation of $\pi^{H_{\Delta,k}}$ and we have $M_k = N_0 \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}} \circ (s^k M)$.

Lemma 4.11. *For any $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ the N_0 -subrepresentation M_k lies in $\mathcal{M}_{\Delta,k}(\pi^{H_{\Delta,k}})$.*

Proof. Let $\{m_1, \dots, m_r\}$ be a set of generators of M as an $A[[N_{\Delta,0}]] [F_\Delta]$ -module. Then by Lemma 4.10 the elements $1 \otimes m_1, \dots, 1 \otimes m_r$ generate $A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}], \varphi_{s^k}]} M$ as a module over $A[[N_{\Delta,k}]] [F_{\Delta,k}]$. In particular, M_k is finitely generated.

For the stability under the action of T_0 note that T_0 normalizes both $H_{\Delta,k}$ and $s^k H_{\Delta,0} s^{-k}$ and the elements in T_0 commute with s .

Since M is admissible as an $N_{\Delta,0}$ -representation and $A[[N_{\Delta,k}]]$ is finitely generated and free as a module over $A[[s^k N_{\Delta,0} s^{-k}]]$, we obtain that M_k is admissible, too. □

In order to simplify notation we write $M_k^\vee[1/X_\Delta] := M_k^\vee[1/\varphi_{s^{kn_0}}(X_\Delta)]$ for $A((N_{\Delta,k})) \otimes_{A[[N_{\Delta,k}]]} M_k^\vee$.

Proposition 4.12. *The map*

$$\begin{aligned} A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M &\rightarrow M_k \\ \lambda \otimes m &\mapsto \lambda \text{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}}(s^k m). \end{aligned} \quad (20)$$

induces an isomorphism $M_k^\vee[1/X_\Delta] \cong A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta]$.

Proof. Since $A[[N_{\Delta,k}]]$ is a finitely generated free module over $A[[s^k N_{\Delta,0} s^{-k}]]$, we have identifications

$$\begin{aligned} (A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M)^\vee &\cong A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M^\vee; \\ (A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M)^\vee[1/X_\Delta] &\cong A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} (M^\vee[1/X_\Delta]). \end{aligned}$$

Therefore we have an injective morphism $f: M_k^\vee[1/X_\Delta] \hookrightarrow A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta]$ of $A((N_{\Delta,k}))$ -modules. Moreover, we have a commutative diagram

$$\begin{array}{ccc} A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M &\twoheadrightarrow & M_k \\ \text{Tr}_{H_{\Delta,0}/H_{\Delta,k}} \downarrow & & \downarrow \text{Tr}_{H_{\Delta,0}/H_{\Delta,k}} \\ A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M &\twoheadrightarrow & M_k \end{array}$$

such that we have $\text{Tr}_{H_{\Delta,0}/H_{\Delta,k}}(M_k) = N_0 F_{s^k}(M) \subseteq M$ and the image of the left vertical map equals $(A[[N_{\Delta,k}]] \otimes_{A[[N_{\Delta,0}],\varphi_{s^k}]} M)^{H_{\Delta,0}}$. Dualizing and inverting $\varphi_{s^{kn_0}}(X_\Delta)$ we obtain a commutative diagram

$$\begin{array}{ccc} A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta] &\xleftarrow{f} \hookrightarrow & M_k^\vee[1/X_\Delta] \\ \text{Tr}_{H_{\Delta,0}/H_{\Delta,k}} \downarrow & & \downarrow \text{Tr}_{H_{\Delta,0}/H_{\Delta,k}} \\ A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta] &\xleftarrow{f} \hookrightarrow & M_k^\vee[1/X_\Delta] \end{array}$$

By Prop. 2.3 the inclusion $N_0 F_{s^k}(M) \subseteq M$ induces an isomorphism

$$M^\vee[1/X_\Delta] \cong A((N_{\Delta,0})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta] = (N_0 F_{s^k}(M))^\vee[1/X_\Delta].$$

On the other hand, the (co)image of the left vertical map is

$$H_0(H_{\Delta,0}/H_{\Delta,k}, A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta]) \cong A((N_{\Delta,0})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta].$$

So f becomes onto after taking $H_{\Delta,0}/H_{\Delta,k}$ -coinvariants. Hence by Nakayama's Lemma f is an isomorphism. \square

Since the map (20) is a $A[[N_{\Delta,k}]] [F_{\Delta,k}]$ -module homomorphism, we obtain a commutative diagram

$$\begin{array}{ccc} A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta] &\xleftarrow{\sim} & M_k^\vee[1/X_\Delta] \\ (1 \otimes F_t)^\vee \downarrow & & \downarrow (1 \otimes F_t)^\vee \\ A((N_{\Delta,k})) \otimes_{A((N_{\Delta,k}),\varphi_t)} A((N_{\Delta,k})) \otimes_{A((N_{\Delta,0}),\varphi_{s^k})} M^\vee[1/X_\Delta] &\xleftarrow{\sim} & A((N_{\Delta,k})) \otimes_{A((N_{\Delta,k}),\varphi_t)} M_k^\vee[1/X_\Delta] \end{array}$$

for all $t \in \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ with both horizontal and the left vertical arrows being isomorphisms. Therefore the right vertical arrow is also an isomorphism. In particular, $M_k^\vee[1/X_\Delta] \cong \mathbb{M}_{k,0}(M^\vee[1/X_\Delta])$ is an étale T_+ -module over $A((N_{\Delta,k}))$ since we have $T_+ = T_0 \prod_{\alpha \in \Delta} t_\alpha^{\mathbb{N}}$ and T_0 acts on $M_k^\vee[1/X_\Delta]$ by conjugation. Taking the projective limit with respect to $k \geq 0$ we define

$$D_{\Delta,\infty}^\vee(\pi) := \varprojlim_{k \geq 0, M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})} M_k^\vee[1/X_\Delta].$$

By construction, $M_\infty^\vee[1/X_\Delta] := \varprojlim_k M_k^\vee[1/X_\Delta]$ is an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,\infty})))$ that corresponds to $M^\vee[1/X_\Delta]$ under the equivalence of categories in Thm. 4.9.

We call two elements $M, M' \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ equivalent ($M \sim M'$) if the inclusions $M \subseteq M + M'$ and $M' \subseteq M + M'$ induce isomorphisms $M^\vee[1/X_\Delta] \cong (M + M')^\vee[1/X_\Delta] \cong M'^\vee[1/X_\Delta]$. In particular, this is an equivalence relation on the set $\mathcal{M}(\pi^{H_{\Delta,0}})$. Similarly, we say that $M_k, M'_k \in \mathcal{M}_{\Delta,k}(\pi^{H_{\Delta,k}})$ are equivalent if the inclusions $M_k \subseteq M_k + M'_k$ and $M'_k \subseteq M_k + M'_k$ induce isomorphisms $M_k^\vee[1/X_\Delta] \cong (M_k + M'_k)^\vee[1/X_\Delta] \cong M'^\vee_k[1/X_\Delta]$.

Remark. The maps

$$\begin{aligned} M &\mapsto N_0 \mathrm{Tr}_{H_{\Delta,k}/s^k H_{\Delta,0} s^{-k}}(s^k M) \\ \mathrm{Tr}_{H_{\Delta,0}/H_{\Delta,k}}(M_k) &\leftarrow M_k \end{aligned}$$

induce a bijection between the sets $\mathcal{M}(\pi^{H_{\Delta,0}})/\sim$ and $\mathcal{M}_{\Delta,k}(\pi^{H_{\Delta,k}})/\sim$. In particular, we have

$$D_{\Delta,\infty}^\vee(\pi) = \varprojlim_{k \geq 0} \varprojlim_{M_k \in \mathcal{M}_{\Delta,k}(\pi^{H_{\Delta,k}})} M_k^\vee[1/X_\Delta].$$

On a finitely generated étale T_+ -module D over $A((N_{\Delta,\infty}))$ we define the weak topology as follows. We put the natural compact topology on any finitely generated $A[[N_{\Delta,k}]]$ -submodule of the $A((N_{\Delta,k}))$ -module $D_{H_{\Delta,k}} = H_0(H_{\Delta,k}, D)$ of coinvariants and the inductive limit topology of these topologies on $D_{H_{\Delta,k}}$ (we call this also the weak topology on $D_{H_{\Delta,k}}$). Now we equip D with the projective limit topology of the weak topologies on $D_{H_{\Delta,k}}$. Finally, if $\varprojlim_{i \in I} D_i$ is a projective limit of finitely generated étale T_+ -modules over $A((N_{\Delta,\infty}))$ then we equip it with the projective limit topology of the weak topologies of each D_i .

4.4 A natural transformation from the Schneider–Vigneras D -functor to $D_{\Delta,\infty}^\vee$

In order to avoid confusion we denote by $D_{SV}(\pi)$ the Schneider–Vigneras functor defined as $D(\pi)$ in [25] (note that in [25] the notation V is used for the \mathfrak{o} -torsion representation that we denote by π). Recall that $D_{SV}(\pi)$ is defined as the inductive limit

$$D_{SV}(\pi) := \varinjlim_{W \in \mathcal{B}_+(\pi)} W^\vee$$

where $\mathcal{B}_+(\pi)$ is the set of generating B_+ -subrepresentations in $\pi|_B$. This is a $\Lambda(N_0)$ -module with an action of B_+^{-1} . For further details we refer the reader to [25]. The inclusions $M_k \subseteq \pi$ ($k \geq 0, M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$) define a T_0 -equivariant $A[[N_0]]$ -module homomorphism $\beta: \pi^\vee \rightarrow$

$D_{\Delta,\infty}^\vee(\pi)$. Our goal is to show that β factors through the map $\pi^\vee \twoheadrightarrow D_{SV}(\pi)$ and it is T_+^{-1} -equivariant. Here T_+^{-1} acts on $D_{\Delta,\infty}^\vee(\pi)$ via the ψ -action (see section 4 of [17]).

Let W be in $\mathcal{B}_+(\pi)$ and $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. Then by Lemma 2.1 in [25] (see also Lemma 3.1 in [17]) there is an integer $k(M) \geq 0$ such that for all $k \geq k(M)$ we have $M_k \leq W$. Therefore the map $\pi^\vee \twoheadrightarrow M_k^\vee \rightarrow M_k^\vee[1/X_\Delta]$ factors through $\pi^\vee \twoheadrightarrow W^\vee$. Taking projective limits with respect to k we obtain a $A[[N_0]]$ -homomorphism

$$\mathrm{pr}_{W,M}: W^\vee \rightarrow M_\infty^\vee[1/X_\Delta].$$

The following is the analogue of Lemma 3.2 in [17] in our situation with essentially the same proof.

Lemma 4.13. *The map $\mathrm{pr}_{W,M}$ is ψ_t -equivariant for all $t \in T_+$.*

Proof. The T_0 -equivariance is clear as it is given by the multiplication by elements of T_0 on both sides. By multiplicativity we are reduced to showing the ψ_t -equivariance for $t = t_\alpha$ for all α in Δ . Let $k = k(\alpha, M) > 0$ be large enough so that $H_{\Delta,k}$ is contained in $t_\alpha H_{\Delta,0} t_\alpha^{-1} \leq t_\alpha N_0 t_\alpha^{-1}$ and M_k is contained in W . Denote by $H_{k,-,\alpha}$ the kernel of the group homomorphism $t_\alpha(\cdot)t_\alpha^{-1}: N_{\Delta,k} \rightarrow N_{\Delta,k}$ and put $\mathrm{Tr} := \mathrm{Tr}_{H_{k,-,\alpha}} = \sum_{u \in H_{k,-,\alpha}} u \in A[[N_{\Delta,k}]]$. By our assumption on k we have $H_{k,-,\alpha} = t_\alpha^{-1} H_{\Delta,k} t_\alpha / H_{\Delta,k}$. For any f in $W^\vee = \mathrm{Hom}_A(W, A)$ and $w \in W$ we have $\psi_{t_\alpha}(f)(w) = f(t_\alpha w)$ by definition. On the other hand, the map $1 \otimes F_{\alpha,k}: A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}]]} M_k \rightarrow M_k$ factors through $A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}/H_{k,-,\alpha}]]} (M_k / \mathrm{Ker}(\mathrm{Tr}|_{M_k}))$. Moreover, $A[[N_{\Delta,k}]]$ is a finite free module over $A[[N_{\Delta,k}/H_{k,-,\alpha}]]$ via φ_{t_α} . Hence we have a series of maps

$$\begin{aligned} M_k^\vee &\xrightarrow{(1 \otimes F_{\alpha,k})^\vee} (A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}/H_{k,-,\alpha}]]} (M_k / \mathrm{Ker}(\mathrm{Tr})))^\vee \xrightarrow{\sim} \\ &\xrightarrow{\sim} A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}/H_{k,-,\alpha}]]} (M_k / \mathrm{Ker}(\mathrm{Tr}))^\vee \xrightarrow{\mathrm{Tr}} \\ &\xrightarrow{\mathrm{Tr}} A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}/H_{k,-,\alpha}]]} \mathrm{Tr}(M_k)^\vee \xrightarrow{\sim} \\ &\xrightarrow{\sim} A[[N_{\Delta,k}]] \otimes_{\varphi_{t_\alpha}, A[[N_{\Delta,k}/H_{k,-,\alpha}]]} (M_k^\vee / \mathrm{Ker}(\mathrm{Tr})) \end{aligned}$$

under which the image of $f = f|_{M_k}$ is as follows:

$$\begin{aligned} &f \xrightarrow{(1 \otimes F_{\alpha,k})^\vee} (f_1: u \otimes (m + \mathrm{Ker}(\mathrm{Tr})) \mapsto f(uF_{\alpha,k}(m))) \xrightarrow{\sim} \\ \xrightarrow{\sim} &\sum_{u \in J(N_{\Delta,k}/t_\alpha N_{\Delta,k} t_\alpha^{-1})} u \otimes f_{3,u} \text{ where } f_{3,u}(m + \mathrm{Ker}(\mathrm{Tr})) := f(uF_{\alpha,k}(m)); \xrightarrow{\sim} \\ &\xrightarrow{\sim} \sum_{u \in J(N_{\Delta,k}/t_\alpha N_{\Delta,k} t_\alpha^{-1})} u \otimes f_{4,u} \text{ where } f_{4,u}(\mathrm{Tr}(m)) := f(ut_\alpha \mathrm{Tr}(m)); \xrightarrow{\sim} \\ &\xrightarrow{\sim} \sum_{u \in J(N_{\Delta,k}/t_\alpha N_{\Delta,k} t_\alpha^{-1})} u \otimes (\mathrm{pr}_{W,M,k}(\psi_{t_\alpha}(u^{-1}f)) + \mathrm{Ker}(\mathrm{Tr})). \end{aligned}$$

Moreover, we have an identification $M_k^\vee[1/X_\Delta] \cong \mathbb{M}_{k,0}(M^\vee[1/X_\Delta])$. In particular, $M_k^\vee[1/X_\Delta]$ is induced as a representation of $H_{k,-,\alpha} \leq H_{\Delta,0}/H_{\Delta,k}$. Therefore $(M_k^\vee / \mathrm{Ker}(\mathrm{Tr}))[1/X_\Delta]$ can be identified with the coinvariants $M_k^\vee[1/X_\Delta]_{H_{k,-,\alpha}}$. So the image of f under the composite map

$$\begin{aligned} W^\vee &\rightarrow M_k^\vee \rightarrow A((N_{\Delta,k})) \otimes_{\varphi_{t_\alpha}, A((N_{\Delta,k}/H_{k,-,\alpha}))} M_k^\vee[1/X_\Delta]_{H_{k,-,\alpha}} \cong \\ &\cong A((N_{\Delta,k})) \otimes_{\varphi_{t_\alpha}, A((N_{\Delta,k}))} M_k^\vee[1/X_\Delta] \end{aligned}$$

equals $\sum_{u \in J(N_{\Delta,k}/t_\alpha N_{\Delta,k} t_\alpha^{-1})} u \otimes \text{pr}_{W,M,k}(\psi_{t_\alpha}(u^{-1}f))$. However, since $M_\infty^\vee[1/X]$ is an étale T_+ -module over $A((N_{\Delta,\infty}))$, we have a unique decomposition of $\text{pr}_{W,M}(f)$ as

$$\text{pr}_{W,M}(f) = \sum_{u \in J(N_0/t_\alpha N_0 t_\alpha^{-1})} u \varphi(\psi_{t_\alpha}(u^{-1} \text{pr}_{W,M}(f)))$$

whence we must have $\psi_{t_\alpha}(\text{pr}_{W,M}(f)) = \text{pr}_{W,M}(\psi_{t_\alpha}(f))$ as claimed. \square

By taking the projective limit with respect to $M \in \mathcal{M}(\pi^{H_{\Delta,0}})$ and the injective limit with respect to $W \in \mathcal{B}_+(\pi)$ we obtain a ψ_s - and Γ -equivariant $\Lambda(N_0)$ -homomorphism

$$\text{pr} := \lim_{\substack{\longrightarrow \\ W}} \lim_{\substack{\longleftarrow \\ M}} \text{pr}_{W,M} : D_{SV}(\pi) \rightarrow D_{\Delta,\infty}^\vee(\pi) .$$

Proposition 4.14. *Let D be a finitely generated étale T_+ -module over $A((N_{\Delta,\infty}))$, and $f : D_{SV}(\pi) \rightarrow D$ be a continuous ψ_t -equivariant $A[[N_0]]$ -homomorphism for all $t \in T_+$. Then f factors uniquely through pr , ie. there exists a unique ψ_t -equivariant $A[[N_0]]$ -homomorphism $\hat{f} : D_{\Delta,\infty}^\vee(\pi) \rightarrow D$ such that $f = \hat{f} \circ \text{pr}$.*

Proof. We prove the uniqueness first. Let \hat{f} and \hat{f}' be two such maps. Then the image of pr lies in the kernel of $\hat{f} - \hat{f}' : D_{\Delta,\infty}^\vee(\pi) \rightarrow D$. By continuity, $\hat{f} - \hat{f}'$ factors through $M_\infty^\vee[1/X_\Delta]$ for some $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. Applying $A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} \cdot$ we deduce that the composite map

$$\pi^\vee \rightarrow M_k^\vee \rightarrow M_k^\vee[1/X_\Delta] = A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} M_\infty^\vee[1/X_\Delta] \xrightarrow{1 \otimes (\hat{f} - \hat{f}')} A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} D$$

is 0. Since the image of M_k^\vee in $M_k^\vee[1/X_\Delta]$ generates $M_k^\vee[1/X_\Delta]$ (as a module over $A((N_{\Delta,k}))$), we deduce that the map $M_k^\vee[1/X_\Delta] \xrightarrow{1 \otimes (\hat{f} - \hat{f}')} A((N_{\Delta,k})) \otimes_{A((N_{\Delta,\infty}))} D$ is 0. Letting $k \rightarrow \infty$ we obtain $\hat{f} = \hat{f}'$ using the remark after Thm. 4.9.

At first we construct a homomorphism $\hat{f}_{H_{\Delta,0}} : D_{\Delta,\infty}^\vee(\pi) = (D_{\Delta,\infty}^\vee(\pi))_{H_{\Delta,0}} \rightarrow D_{H_{\Delta,0}}$ such that the following diagram commutes:

$$\begin{array}{ccccc} D_{SV}(\pi) & \xrightarrow{\text{pr}} & D_{\Delta,\infty}^\vee(\pi) & \xrightarrow{(\cdot)_{H_{\Delta,0}}} & D_{\Delta}^\vee(\pi) \\ & \searrow f & & & \downarrow \hat{f}_{H_{\Delta,0}} \\ & & D & \xrightarrow{(\cdot)_{H_{\Delta,0}}} & D_{H_{\Delta,0}} \end{array}$$

Consider the composite map $f' : \pi^\vee \rightarrow D_{SV}(\pi) \xrightarrow{f} D \rightarrow D_{H_{\Delta,0}}$. Note that f' is continuous and $D_{H_{\Delta,0}}$ is Hausdorff, so $\text{Ker}(f')$ is closed in π^\vee . Therefore $M_0 := (\pi^\vee / \text{Ker}(f'))^\vee$ is naturally a subspace in π . We claim that M_0 lies in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. Indeed, M_0^\vee is a quotient of $\pi_{H_{\Delta,0}}^\vee$, hence $M_0 \leq \pi^{H_{\Delta,0}}$ and it is T_0 -invariant since f' is T_0 -equivariant. M_0 is admissible because it is discrete, hence M_0^\vee is compact, equivalently finitely generated over $A[[N_{\Delta,0}]]$, because M_0^\vee can be identified with a $A[[N_{\Delta,0}]]$ -submodule of $D_{H_{\Delta,0}}$ which is finitely generated over $A((N_{\Delta,0}))$. Finally, M_0 is finitely generated over $A[[N_{\Delta,0}]][[F_\Delta]]$ by Proposition 2.8.

Now we have an injective morphism $f_0 : M_0^\vee[1/X_\Delta] \hookrightarrow D$ of étale T_+ -modules over $A((N_{\Delta,0}))$. The map $\hat{f}_{H_{\Delta,0}} : D_{\Delta}^\vee(\pi) \rightarrow D_{H_{\Delta,0}}$ is the composite map $D_{\Delta}^\vee(\pi) \rightarrow M_0^\vee[1/X_\Delta] \hookrightarrow D$. It is well defined and makes the above diagram commutative, because the map

$$\pi^\vee \rightarrow D_{SV}(\pi) \xrightarrow{\text{pr}} D_{\Delta,\infty}^\vee(\pi) \xrightarrow{(\cdot)_{H_{\Delta,0}}} D_{\Delta}^\vee(\pi) \rightarrow M_0^\vee[1/X_\Delta]$$

is the same as $\pi^\vee \rightarrow M_0^\vee \rightarrow M_0^\vee[1/X_\Delta]$.

Finally, $M^\vee[1/X]$ (resp. $D_{H_{\Delta,0}}$) corresponds to $M_\infty^\vee[1/X_\Delta]$ (resp. to D) via the equivalence of categories in Theorem 4.9 therefore f_0 can uniquely be lifted to a T_+ -equivariant $A((N_{\Delta,\infty}))$ -homomorphism $f_\infty: M_\infty^\vee[1/X_\Delta] \hookrightarrow D$. The map \hat{f} is defined as the composite $D_{\Delta,\infty}^\vee(\pi) \rightarrow M_\infty^\vee[1/X_\Delta] \hookrightarrow D$. Now the image of $f - \hat{f} \circ \text{pr}$ is a ψ_t -invariant ($t \in T_+$) $A[[N_0]]$ -submodule in $(H_{\Delta,0} - 1)D$. For any $x \in D_{SV}(\pi)$ and $k \geq 0$ we may write $(f - \hat{f} \circ \text{pr})(x)$ in the form $\sum_{u \in J(N_0/s^k N_0 s^{-k})} u \varphi^k((f - \hat{f} \circ \text{pr})(\psi^k(u^{-1}x)))$ that lies in $(H_{\Delta,k} - 1)D$. Hence we obtain $(f - \hat{f} \circ \text{pr})(D_{SV}(\pi)) \subseteq \bigcap_k (H_{\Delta,k} - 1)D = \{0\}$ as we have $D \cong \mathbb{M}_{\infty,0}(D_{H_{\Delta,0}}) = \varprojlim_k D_{H_{\Delta,k}}$. \square

By Corollary 4.11 in [17] there exists a homomorphism $\tilde{\text{pr}}: \widetilde{D_{SV}(\pi)} \rightarrow D_{\Delta,\infty}^\vee(\pi)$ of étale T_+ -modules over $A[[N_0]]$ such that $\text{pr} = \tilde{\text{pr}} \circ \iota$ where $\iota: D_{SV}(\pi) \hookrightarrow \widetilde{D_{SV}(\pi)}$ is the étale hull of $D_{SV}(\pi)$ which is defined as $\widetilde{D_{SV}(\pi)} := \varinjlim_{t \in T_+} \varphi_t^* D_{SV}(\pi)$ where we put $\varphi_t^* D_{SV}(\pi) := A[[N_0]]_{A[[N_0]], \varphi_t} D_{SV}(\pi)$. For more details on the étale hull, in particular its universal property, we refer the reader to [17]. The following is the natural analogue of Thm. 2.27 in [17] in our situation.

Corollary 4.15. *We have*

$$D_{\Delta,\infty}^\vee(\pi) \cong \varprojlim_D D$$

where D runs through the finitely generated étale T_+ -modules over $A((N_{\Delta,\infty}))$ arising as a quotient of $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D_{SV}(\pi)}$ such that the quotient map is continuous in the weak topology of D and the final topology on $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D_{SV}(\pi)}$ of the map $1 \otimes \iota: D_{SV}(\pi) \rightarrow A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D_{SV}(\pi)}$.

Proof. Let D be a finitely generated étale T_+ -module over $A((N_{\Delta,\infty}))$ with a continuous ψ_t -equivariant (for all $t \in T_+$) homomorphism of $A[[N_0]]$ -modules $D_{SV}(\pi) \rightarrow D$. Then by the universal property (Prop. 2.20 in [17]) of the étale hull this factors uniquely through $\widetilde{D_{SV}(\pi)}$ and also through $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D_{SV}(\pi)}$ since D is a module over $A((N_{\Delta,\infty}))$. Therefore by Lemma 4.13 and Proposition 4.14 the topological quotients of $A((N_{\Delta,\infty})) \otimes_{A[[N_0]]} \widetilde{D_{SV}(\pi)}$ in the category of finitely generated étale T_+ -modules over $A((N_{\Delta,\infty}))$ are exactly the étale T_+ modules $M_\infty^\vee[1/X_\Delta]$ for $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. \square

4.5 A G -equivariant sheaf on G/B attached to $D_\Delta^\vee(\pi)$

The goal of this section is to construct a G -equivariant sheaf on G/B with sections on $\mathcal{C}_0 := N_0 w_0 B/B$ isomorphic to $\widetilde{D_\Delta(\pi)} := \tilde{\text{pr}}(\widetilde{D_{SV}(\pi)}) \subseteq D_{\Delta,\infty}^\vee(\pi)$ as an étale T_+ -module. (Here w_0 denotes a representative of the element with maximal length in the Weyl group $N_G(T)/T$.) The method of constructing a G -equivariant sheaf on G/B does not work in our situation since the property $\mathfrak{T}(1)$ in Prop. 6.8 in [26] is not satisfied in general for étale T_+ -modules over $A((N_{\Delta,\infty}))$. The idea is to use the G -action on π^\vee in order to construct the operators $\mathcal{H}_g: \widetilde{D_\Delta(\pi)} \rightarrow \widetilde{D_\Delta(\pi)}$. So let π be a smooth representation of G over A and choose an arbitrary object M in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. Denote by M_∞ the Pontryagin dual of the image of the natural map $\pi^\vee \rightarrow M_\infty^\vee[1/X_\Delta]$.

Lemma 4.16. M_∞ is a B_+ -subrepresentation of π . If M is chosen so that M^\vee is X_Δ -torsion free then we have $M_\infty = \bigcup_k M_k = B_+M$.

Remark. Let D_0 be the image of π^\vee in $M^\vee[1/X_\Delta]$. By Proposition 2.8 $D_0^\vee \subseteq M$ lies in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ and $D_0 \cong (D_0^\vee)^\vee$ is X_Δ -torsion free since it is contained in $M^\vee[X_\Delta]$. Moreover, we have $D_0[1/X_\Delta] = M^\vee[1/X_\Delta]$, so we may replace M by D_0^\vee so that the second conclusion holds.

Proof. The N_0 -invariance of the subspace $M_\infty \subseteq \pi$ is clear. Let $t \in T_+$ and $m \in M_\infty$ be arbitrary. Assume that tm does not lie in $M_\infty \leq \pi$. Then there is an element $\mu \in \pi^\vee$ such that $\mu|_{M_\infty} = 0$ but $(t^{-1}\mu)(m) = \mu(tm) \neq 0$. By Lemma 4.13 we compute $0 \neq (t^{-1}\mu)|_{M_\infty} = \text{pr}_{W,M}(t^{-1}\mu|_W) = \psi_t(\text{pr}_{W,M}(\mu)) = \psi_t(0) = 0$ which is a contradiction. So M_∞ is B_+ -invariant.

Assume now that M^\vee is X_Δ -torsion free, ie. the map $M^\vee \rightarrow M^\vee[1/X_\Delta]$ is injective. Therefore M is contained in M_∞ since M^\vee is a quotient of M_∞^\vee . Hence $M_k \subseteq B_+M$ is also contained in M_∞ . Now assume that $\bigcup_k M_k \subsetneq M_\infty$. Then there exists an element μ in π^\vee such that $\mu|_{M_k} = 0$ for all $k \geq 0$ but $\mu|_{M_\infty} \neq 0$. In particular, the image of μ in $M_k^\vee[1/X_\Delta]$ is zero for all $k \geq 0$ whence it is also zero in $\varprojlim M_k^\vee[1/X_\Delta] = M_\infty^\vee[1/X_\Delta]$ contradicting to $\mu|_{M_\infty} \neq 0$. \square

Let us denote by $\mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$ the set of those $M \in \mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ so that M^\vee has no X_Δ -torsion. We still have $D_\Delta^\vee(\pi) = \varprojlim_{M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})} M^\vee[1/X_\Delta]$.

Lemma 4.17. Suppose we choose $M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$. Then the map

$$1 \otimes F_s : A[[N_{\Delta,0}]] \otimes_{\varphi_s, A[[N_{\Delta,0}]]} M \rightarrow M$$

is surjective.

Proof. The assertion is equivalent to the injectivity of the dual map

$$(1 \otimes F_s)^\vee : M^\vee \rightarrow A[[N_{\Delta,0}]] \otimes_{\varphi_s, A[[N_{\Delta,0}]]} M^\vee .$$

This map is, however, injective by Lemma 4.2, Prop. 4.5, and Cor. 4.8 in [17] noting that M^\vee is contained in the étale T_+ -module $M^\vee[1/X_\Delta]$. \square

For an integer $n \geq 1$ we denote by $U^{(n)}$ the kernel of the group homomorphism $\mathbf{G}(\mathbb{Z}_p) \rightarrow \mathbf{G}(\mathbb{Z}/p^n\mathbb{Z})$.

Lemma 4.18. We have $U^{(1)}M_\infty = M_\infty$ for all $M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$.

Proof. We may assume without loss of generality that M^\vee is X_Δ -torsion free so that we have $M_\infty = B_+M$ by Lemma 4.16. Now since M lies in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$, it is generated by finitely many elements $m_1, \dots, m_r \in M$ as a module over $A[[N_{\Delta,0}]] [F_\Delta]$. Since the elements of $A[[N_{\Delta,0}]] [F_\Delta]$ are finite linear combinations of elements in B_+ we deduce that m_1, \dots, m_r generates M_∞ as a B_+ -subrepresentation of π . Since π is smooth, there exists an integer $n \geq 1$ such that m_1, \dots, m_r are all fixed by $U^{(n)}$. Moreover, we may assume without loss of generality that the group T_0 permutes the elements of the set $\{m_1, \dots, m_r\}$. By Lemmata 4.16 and 4.17 we may write any element $m \in M_\infty$ as a finite linear combination $m = \sum_{i=1}^r \lambda_i s^n m_i$ with λ_i in the monoid-ring $A[B_+]$. So we are reduced to showing that $uvs^n m_i$ lies in M_∞ for all

$u \in U^{(1)}$, $v \in N_0$, $t \in T_+$, and $1 \leq i \leq r$. Since $U^{(1)}$ is normal in $\mathbf{G}(\mathbb{Z}_p)$ and $N_0 \subseteq \mathbf{G}(\mathbb{Z}_p)$, the element $u_1 := v^{-1}uv$ also lies in $U^{(1)}$. By the Iwahori factorization we may write u_1 as a product $u_1 = n_1 t_1 \bar{n}_1$ with $n_1 \in U^{(1)} \cap N \leq N_0$, $t_1 \in U^{(1)} \cap T$, and $\bar{n}_1 \in U^{(1)} \cap \bar{N}$. Therefore $s^{-n} t^{-1} \bar{n}_1 t s^n$ lies in $s^{-n} t^{-1} (U^{(1)} \cap \bar{N}) t s^n \leq U^{(n)} \cap \bar{N}$ whence we have $s^{-n} t^{-1} \bar{n}_1 t s^n m_i = m_i$ for all $1 \leq i \leq r$ by our choice of n . So we compute

$$u w t s^n m_i = v u_1 t s^n m_i = v n_1 t_1 t s^n (s^{-n} t^{-1} \bar{n}_1 t s^n) m_i = v n_1 t_1 t s^n m_i \in B_+ M = M_\infty .$$

□

Now let g be an arbitrary element in G and put $\mathcal{U}_g := \{u \in N_0 \mid x_u \in g^{-1} \mathcal{C}_0 \cap \mathcal{C}_0\}$ where $u \mapsto x_u = u w_0 B$ is the homeomorphism $N_0 \rightarrow \mathcal{C}_0 = N_0 w_0 B / B \subset G/B$. For $u \in \mathcal{U}_g$ we may factorize gu as $gu = n(g, u) t(g, u) \bar{n}(g, u)$ with $n(g, u) \in N_0$, $t(g, u) \in T$, and $\bar{n}(g, u) \in \bar{N}$. By Lemma 7.3 in [17] there exists an integer $k_0 = k_0(g)$ such that for all $k \geq k_0$ and $u \in \mathcal{U}_g$ we have $u s^k N_0 s^{-k} \subseteq \mathcal{U}_g$, $s^k t(g, u) \in T_+$, and $s^{-k} \bar{n}(g, u) s^k \in U^{(1)} \cap \bar{N}$. Moreover, since \mathcal{U}_g is compact, it is a finite union of cosets of the form $u s^k N_0 s^{-k}$ ($u \in \mathcal{U}_g$). We denote by $\widetilde{M}_\infty^\vee$ the étale hull of the image $M_\infty^\vee \subset M_\infty^\vee[1/X_\Delta]$ of the natural map $\text{pr}_M: \pi^\vee \rightarrow M_\infty^\vee[1/X_\Delta]$. As a special case of Cor. 4.8 in [17] we may view $\widetilde{M}_\infty^\vee$ as an étale T_+ -submodule of $M_\infty^\vee[1/X_\Delta]$. More explicitly, since the powers of s are cofinal in T_+ , we have

$$\begin{aligned} \widetilde{M}_\infty^\vee &= \bigcup_{k \geq 0} \bigoplus_{u \in J(N_0/s^k N_0 s^{-k})} u \varphi_{s^k}(M_\infty^\vee) = \bigcup_{k \geq k_0} \bigoplus_{u \in J(N_0/s^k N_0 s^{-k})} u \varphi_{s^k}(M_\infty^\vee) = \\ &= \bigcup_{k \geq k_0} \bigoplus_{u \in J(N_0/s^k N_0 s^{-k})} \text{res}_{u s^k N_0 s^{-k}}^{N_0}(M_\infty^\vee) \end{aligned}$$

by Cor. 4.11 in [17] noting that $\psi_s: M_\infty^\vee \rightarrow M_\infty^\vee$ is surjective since $(s \cdot): M_\infty \rightarrow M_\infty$ is injective. Here $\text{res}_{u s^k N_0 s^{-k}}^{N_0}$ denotes the operator $(u \cdot) \circ \varphi_{s^k} \circ \psi_{s^k} \circ (u^{-1} \cdot)$. More generally, for an open compact subset $\mathcal{U} \subset N_0$ we may write \mathcal{U} as a disjoint union of cosets $u s^k N_0 s^{-k}$ (u running on a finite subset $U_k \subset \mathcal{U}$) for $k \geq 0$ large enough depending on \mathcal{U} and put $\text{res}_{\mathcal{U}}^{N_0} := \sum_{u \in U_k} \text{res}_{u s^k N_0 s^{-k}}^{N_0}$. For $k \geq k_0$ we choose a set $U_{k,g} := \mathcal{U}_g \cap J(N_0/s^k N_0 s^{-k})$ of representatives of the cosets of $\mathcal{U}_g/s^k N_0 s^{-k}$. By Lemma 4.13 in [17] the map

$$n(g, \cdot): u s^k N_0 s^{-k} \rightarrow n(g, u) t(g, u) s^k N_0 s^{-k} t(g, u)^{-1}$$

is a bijection for each $u \in U_{k,g}$. Under the identification $N_0 \xrightarrow{\sim} \mathcal{C}_0$ the above bijection corresponds to the bijection

$$(g \cdot): u s^k N_0 w_0 B / B \xrightarrow{\sim} n(g, u) t(g, u) s^k N_0 w_0 B / B .$$

So we define

$$\begin{aligned} \mathcal{H}_g: \widetilde{M}_\infty^\vee &\rightarrow \widetilde{M}_\infty^\vee \\ \sum_{u \in J(N_0/s^k N_0 s^{-k})} \text{res}_{u s^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k})) &\mapsto \sum_{u \in U_{k,g}} \text{res}_{n(g,u)t(g,u)s^k N_0 s^{-k} t(g,u)^{-1}}^{N_0}(\text{pr}_M(g \mu_{u,k})) \end{aligned}$$

for $\mu_{u,k}$ in π^\vee ($u \in U_{k,g}$, $k \geq k_0$).

Lemma 4.19. *The map \mathcal{H}_g above is well-defined.*

Proof. Since $\text{res}_{\mathcal{U}}^{N_0}$ does not depend on the choice of the decomposition of the compact open subset $\mathcal{U} \subseteq N_0$ into a disjoint union of cosets of subgroups of N_0 of the form tN_0t^{-1} ($t \in T_+$), it suffices to show that \mathcal{H}_g does not depend on the choice of $\mu_{u,k}$ ($u \in U_{k,g}$, $k \geq k_0$). Let μ be in π^\vee such that $\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu)) = 0$. Since $(u \cdot) \circ \varphi_{s^k}$ is injective on $\widetilde{M}_\infty^\vee$, we obtain $\text{pr}_M(s^{-k}u^{-1}\mu) = \psi_{s^k}(u^{-1}\text{pr}_M(\mu)) = 0$ by Lemma 4.13. In other words the restriction of $s^{-k}u^{-1}\mu$ to $M_\infty \subset \pi$ is zero. By our assumption $k \geq k_0$ we have $s^{-k}\overline{n}(g,u)s^k$ lies in $U^{(1)}$. Hence by Lemma 4.18 we compute

$$0 = \text{pr}_M(s^{-k}u^{-1}\mu) = \text{pr}(s^{-k}\overline{n}(g,u)u^{-1}\mu) = \text{pr}(s^{-k}t(g,u)^{-1}n(g,u)^{-1}g\mu) .$$

In particular, the formula defining \mathcal{H}_g does not depend on the choice of $\mu_{u,k}$. \square

Proposition 4.20. *For any smooth o -torsion representation π of G and any $M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$ there exists a G -equivariant sheaf $\mathfrak{Y}_{\pi,M}$ on G/B with sections $\mathfrak{Y}_{\pi,M}(\mathcal{C}_0)$ on \mathcal{C}_0 isomorphic to $\widetilde{M}_\infty^\vee$ as an étale T_+ -module over $A[[N_0]]$ such that we have $\mathcal{H}_g = (g \cdot) \circ \text{res}_{g^{-1}\mathcal{C}_0 \cap \mathcal{C}_0}^{\mathcal{C}_0}$ as maps on $\mathfrak{Y}_{\pi,M}(\mathcal{C}_0)$.*

Proof. By Prop. 5.14 in [26] we are bound to check the relations H1, H2, H3 therein: for any $g, h \in N_0\overline{B}N_0$, $b \in B \cap N_0\overline{B}N_0$ and compact open subset $\mathcal{V} \subset \mathcal{C}_0$ we have

$$\text{H1 } \text{res}_{\mathcal{V}}^{\mathcal{C}_0} \circ \mathcal{H}_g = \mathcal{H}_g \circ \text{res}_{g^{-1}\mathcal{V} \cap \mathcal{C}_0}^{\mathcal{C}_0} ;$$

$$\text{H2 } \mathcal{H}_g \circ \mathcal{H}_h = \mathcal{H}_{gh} \circ \text{res}_{(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0}^{\mathcal{C}_0} ;$$

$$\text{H3 } \mathcal{H}_b = b \circ \text{res}_{b^{-1}\mathcal{C}_0 \cap \mathcal{C}_0}^{\mathcal{C}_0} .$$

Let d be in $\widetilde{M}_\infty^\vee$ and let $k \geq \max(k_0(g), k_0(h), k_0(gh))$ be large enough so that all the sets $g^{-1}\mathcal{V} \cap \mathcal{C}_0$, $\mathcal{U}_g = g^{-1}\mathcal{C}_0 \cap \mathcal{C}_0$, $(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0$, \mathcal{U}_h , and \mathcal{U}_b can be written as the disjoint union of sets of the form $us^kN_0w_0B/B$ for $u \in N_0$ and so that d is contained in $N_0\varphi^k(M_\infty^\vee)$. So we may write $d = \sum_{u \in J(N_0/s^kN_0s^{-k})} \text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))$ for some $\mu_{u,k} \in \pi^\vee$ ($u \in J(N_0/s^kN_0s^{-k})$).

H1: We distinguish two cases: If x_u does not lie in $g^{-1}\mathcal{V} \cap \mathcal{C}_0$ then we have $us^kN_0w_0B \cap g^{-1}\mathcal{V} = \emptyset$ whence the right hand side of H1 has value zero at $\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))$. Moreover, we obtain $n(g,u)t(g,u)s^kN_0w_0B \cap \mathcal{V} = gus^kN_0w_0B \cap \mathcal{V} = \emptyset$ (Lemma 7.4 in [17]), so we the left hand side of H1 also has value zero at $\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))$. On the other hand, if x_u lies in $g^{-1}\mathcal{V} \cap \mathcal{C}_0$ then we have $us^kN_0w_0B \subseteq g^{-1}\mathcal{V} \cap \mathcal{C}_0$ and $n(g,u)t(g,u)s^kN_0w_0B \subseteq \mathcal{V} \cap g\mathcal{C}_0$ whence both sides have value $\mathcal{H}_g(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k})))$ at $\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))$.

H2: We distinguish two cases: If x_u does not lie in $(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0$ then we clearly have

$$\mathcal{H}_{gh} \circ \text{res}_{(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0}^{\mathcal{C}_0}(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) = 0 .$$

Further, $\mathcal{H}_h(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k})))$ is supported on $hus^kN_0w_0B$ that is disjoint to \mathcal{U}_g whence we have

$$\mathcal{H}_g \circ \mathcal{H}_h(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) = 0 ,$$

too. On the other hand, if x_u lies in $(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0$ then we compute

$$\begin{aligned} & \mathcal{H}_{gh} \circ \text{res}_{(gh)^{-1}\mathcal{C}_0 \cap h^{-1}\mathcal{C}_0 \cap \mathcal{C}_0}^{\mathcal{C}_0}(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) = \\ & = \mathcal{H}_{gh}(\text{res}_{us^kN_0s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) = \text{res}_{ghus^kN_0w_0B}^{\mathcal{C}_0}(\text{pr}_M(gh\mu_{u,k})) . \end{aligned}$$

Moreover, we write $hus^k N_0 w_0 B$ as the disjoint union of sets of the form $vs^l N_0 w_0 B$ (v running on a subset $V \subseteq J(N_0/s^l N_0 s^{-l})$) for some $l \geq k$. So we compute

$$\begin{aligned} \mathcal{H}_g \circ \mathcal{H}_h(\text{res}_{us^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) &= \mathcal{H}_g \circ \text{res}_{hus^k N_0 w_0 B}^{C_0}(\text{pr}_M(h\mu_{u,k})) = \\ &= \sum_{v \in V} \mathcal{H}_g \circ \text{res}_{vs^l N_0 w_0 B}^{C_0}(\text{pr}_M(h\mu_{u,k})) = \sum_{v \in V} \text{res}_{gvs^l N_0 w_0 B}^{C_0}(\text{pr}_M(gh\mu_{u,k})) = \\ &= \text{res}_{ghus^k N_0 w_0 B}^{C_0}(\text{pr}_M(gh\mu_{u,k})) . \end{aligned}$$

H3: Let $b = v_1 t v_2$ be in $B \cap N_0 \overline{B} N_0 = N_0 T N_0$ with $v_1, v_2 \in N_0$ and $t \in T$. We may write $t = t_1^{-1} t_2$ for some $t_1, t_2 \in T_+$. Recall from [26] that the action of b on $\widetilde{M}_\infty^\vee$ is defined by the formula $b(d) = v_1 \psi_{t_1} \circ \varphi_{t_2}(v_2 d)$. This does not depend on the choice of t_1 and t_2 as we have $\psi_{t'} \circ \varphi_{t'} = \text{id}$ for all $t' \in T_+$. If x_u does not lie in $b^{-1} \mathcal{C}_0 \cap \mathcal{C}_0$ then both sides of H3 vanish at $\text{res}_{us^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))$, so assume we have $x_u \in b^{-1} \mathcal{C}_0 \cap \mathcal{C}_0$. By the choice of k and u , ts^k lies in T_+ and $tv_2 ut^{-1}$ lies in N_0 . We compute

$$\begin{aligned} \mathcal{H}_b(\text{res}_{us^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) &= \text{res}_{(v_1 t v_2 ut^{-1})ts^k N_0 w_0 B}^{C_0}(\text{pr}_M(v_1 t v_2 \mu_{u,k})) = \\ &= v_1 t v_2 ut^{-1} \varphi_{ts^k} \circ \psi_{ts^k}(tu^{-1} v_2^{-1} t^{-1} v_1^{-1} \text{pr}_M(v_1 t v_2 \mu_{u,k})) = \\ &= v_1 t v_2 ut^{-1} \psi_{t_1} \circ \varphi_{t_2 s^k}(\text{pr}_M(s^{-k} t^{-1} t u^{-1} v_2^{-1} t^{-1} v_1^{-1} v_1 t v_2 \mu_{u,k})) = \\ &= v_1 \psi_{t_1}(t_1 t v_2 ut^{-1} t_1^{-1} \varphi_{t_2} \circ \varphi_{s^k}(\text{pr}_M(s^{-k} u^{-1} \mu_{u,k}))) = \\ &= v_1 \psi_{t_1} \circ \varphi_{t_2}(v_2 u \varphi_{s^k} \circ \psi_{s^k}(u^{-1} \text{pr}_M(\mu_{u,k}))) = \\ &= v_1 \psi_{t_1} \circ \varphi_{t_2}(v_2 \text{res}_{us^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k}))) = b \circ \text{res}_{us^k N_0 s^{-k}}^{N_0}(\text{pr}_M(\mu_{u,k})) \end{aligned}$$

as desired. \square

We equip the space $\mathfrak{Y}_{\pi, M}(G/B)$ of global sections with the coarsest topology such that the restriction maps $\mathfrak{Y}_{\pi, M}(G/B) \rightarrow \mathfrak{Y}_{\pi, M}(g\mathcal{C}_0)$ are continuous for all $g \in G$. This topology is Hausdorff by Lemma 4.9 in [17].

We denote by $\beta_{\mathcal{C}_0, M}$ the composite map $\pi^\vee \rightarrow M_\infty^\vee \rightarrow \widetilde{M}_\infty^\vee = \mathfrak{Y}_{\pi, M}(\mathcal{C}_0)$. For each $g \in G$ and $\mu \in \pi^\vee$ we define $\beta_{g\mathcal{C}_0, M}(\mu) := g \cdot \beta_{\mathcal{C}_0, M}(g^{-1} \mu)$. Writing $g^{-1} \mathcal{C}_0 \cap \mathcal{C}_0$ as a disjoint union of sets of the form $us^k N_0 w_0 B$ with $u \in U$ for some $k \geq k_0(g)$ and $U := J(N_0/s^k N_0 s^{-k}) \cap \mathcal{U}_g$, we compute

$$\begin{aligned} \text{res}_{\mathcal{C}_0 \cap g\mathcal{C}_0}^{g\mathcal{C}_0}(\beta_{g\mathcal{C}_0, M}(\mu)) &= \text{res}_{\mathcal{C}_0 \cap g\mathcal{C}_0}^{g\mathcal{C}_0}(g\beta_{\mathcal{C}_0, M}(g^{-1} \mu)) = g \text{res}_{g^{-1} \mathcal{C}_0 \cap \mathcal{C}_0}^{C_0}(\beta_{\mathcal{C}_0, M}(g^{-1} \mu)) = \\ &= \mathcal{H}_g(\text{pr}_M(g^{-1} \mu)) = \sum_{u \in U} \text{res}_{gus^k N_0 w_0 B}^{C_0}(\text{pr}_M(\mu)) = \\ &= \text{res}_{\mathcal{C}_0 \cap g\mathcal{C}_0}^{C_0}(\text{pr}_M(\mu)) = \text{res}_{\mathcal{C}_0 \cap g\mathcal{C}_0}^{C_0}(\beta_{\mathcal{C}_0, M}(\mu)) . \end{aligned}$$

Since the compact open subsets $g\mathcal{C}_0$ ($g \in G$) cover G/B , the maps $\beta_{g\mathcal{C}_0, M}: \pi^\vee \rightarrow \mathfrak{Y}_{\pi, M}(g\mathcal{C}_0)$ glue together into a continuous G -equivariant map $\beta_{G/B, M}: \pi^\vee \rightarrow \mathfrak{Y}_{\pi, M}(G/B)$ with $\beta_{g\mathcal{C}_0, M} = \text{res}_{g\mathcal{C}_0}^{G/B} \circ \beta_{G/B, M}$ for all $g \in G$.

Corollary 4.21. *The map $\beta_{G/B, M}: \pi^\vee \rightarrow \mathfrak{Y}_{\pi, M}(G/B)$ is natural in the pair (π, M) in the following sense: for a morphism $f: \pi \rightarrow \pi'$ of smooth representations of G and $M \in \mathcal{M}_\Delta^0(\pi^{H\Delta, 0})$,*

the image $f(M)$ lies in $\mathcal{M}_\Delta^0(\pi'^{H_{\Delta,0}})$. Moreover, for any $M' \in \mathcal{M}_\Delta^0(\pi'^{H_{\Delta,0}})$ containing $f(M)$ we have a commutative square

$$\begin{array}{ccc} \pi^\vee & \xrightarrow{\beta_{G/B,M'}} & \mathfrak{Y}_{\pi',M'}(G/B) \\ \downarrow & & \downarrow \\ \pi^\vee & \xrightarrow{\beta_{G/B,M}} & \mathfrak{Y}_{\pi,M}(G/B) \end{array}$$

Now we fix π and vary M in $\mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$ to obtain

Corollary 4.22. *There exists a G -equivariant sheaf $\mathfrak{Y}_{\pi,\Delta}$ on G/B with sections $\mathfrak{Y}_{\pi,\Delta}(\mathcal{C}_0)$ isomorphic to $\widetilde{D}_\Delta(\pi) = \widetilde{\text{pr}}(\widetilde{D}_{SV}(\pi))$ together with a natural G -equivariant continuous map $\beta_{G/B,\Delta}: \pi^\vee \rightarrow \mathfrak{Y}_{\pi,\Delta}(G/B)$.*

Proof. Note that the image of π^\vee in $D_\Delta^\vee(\pi)$ is the Pontryagin dual of the union $\bigcup_{M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})} M_\infty$. Therefore by definition $\widetilde{D}_\Delta(\pi)$ is the étale hull of $(\bigcup_{M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})} M_\infty)^\vee = \varprojlim_{M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})} M_\infty^\vee$. So we may conclude the existence of the operators \mathcal{H}_g as in Lemma 4.19. The existence of the sheaf follows the same way as in Prop. 4.20. \square

Remark. Let π be an irreducible admissible smooth representation of G over κ and assume that $D_\Delta^\vee(\pi) \neq 0$. Then we can realize π^\vee as a G -invariant subspace in the global sections of a “small” G -equivariant sheaf \mathfrak{Y} on G/B . By small we mean here that the section $\mathfrak{Y}(\mathcal{C}_0)$ is contained in a *finitely generated* module over $\kappa((N_{\Delta,\infty}))$. Indeed, we may take any $0 \neq M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$ and put $\mathfrak{Y} := \mathfrak{Y}_{\pi,M}$. It would be natural to expect that the irreducibility of π would imply the irreducibility of $D_\Delta^\vee(\pi)$. In particular, this would mean that $D_\Delta^\vee(\pi)$ is in fact finitely generated. We end this section by presenting some very preliminary ideas towards proving this.

We choose a total ordering of the Weyl group $N_G(T)/T$ refining the Bruhat order. This gives a decreasing filtration of G/B by open B -invariant subsets $\text{Fil}^w(G/B) := \bigcup_{w_1 \geq w} Bw_1B/B \subseteq G/B$ for $w \in N_G(T)/T$. Its bottom term is $\text{Fil}^{w_0}(G/B)$ for the element $w_0 \in N_G(T)/T$ of maximal length and we have $\text{Fil}^1(G/B) = G/B$. For each $w \in N_G(T)/T$ we define

$$\text{Fil}_M^w(\pi^\vee) := \beta_{G/B,M}^{-1}(\text{Ker}(\text{res}_{\text{Fil}^w(G/B)}^{G/B}: \mathfrak{Y}_{\pi,M}(G/B) \rightarrow \mathfrak{Y}_{\pi,M}(\text{Fil}^w(G/B)))) .$$

This is an increasing filtration of π^\vee by closed B -subrepresentations. Taking Pontryagin duals we obtain a decreasing filtration $\text{Fil}_M^w(\pi) := (\pi^\vee / \text{Fil}_M^w(\pi^\vee))^\vee \leq \pi$ whose graded pieces we denote by $\text{gr}_M^w(\pi)$.

Lemma 4.23. *We have $\text{gr}_M^{w_0}(\pi) = \bigcup_{n \geq 0} s^{-n} M_\infty$ and $D_{SV}(\text{gr}_M^{w_0}(\pi)) \cong M_\infty^\vee$.*

Proof. Note that M_∞^\vee can be identified with $\text{res}_{\mathcal{C}_0}^{G/B}(\beta_{G/B,M}(\pi^\vee))$. On the other hand, we write $\mathcal{C} := Nw_0B/B = \text{Fil}^{w_0}(G/B) = \bigcup_{n \geq 0} s^{-n}\mathcal{C}_0$. So we compute

$$\begin{aligned} \text{gr}_M^{w_0}(\pi)^\vee &= \text{res}_{\text{Fil}^{w_0}}^{G/B}(\beta_{G/B,M}(\pi^\vee)) = \varprojlim_n \text{res}_{s^{-n}\mathcal{C}_0}^{G/B} \circ \beta_{G/B,M}(\pi^\vee) = \varprojlim_n \beta_{s^{-n}\mathcal{C}_0,M}(\pi^\vee) = \\ &= \varprojlim_n s^{-n} \cdot \beta_{\mathcal{C}_0,M}(\pi^\vee) = \varprojlim_{\psi_s: M_\infty^\vee \rightarrow M_\infty^\vee} M_\infty^\vee = \left(\varinjlim_{s^n: M_\infty \rightarrow M_\infty} M_\infty \right)^\vee = \left(\bigcup_n s^{-n} M_\infty \right)^\vee . \end{aligned}$$

This shows, in particular, that M_∞ is a generating B_+ -subrepresentation of $\mathrm{gr}_M^{w_0}(\pi)$. Let $W \subseteq M_\infty$ be another generating B_+ -subrepresentation. By Lemma 4.16 the B_+ -subrepresentation M_∞ is generated by a finite set $\{m_1, \dots, m_r\}$ of elements in M . By Lemma 2.1 in [25] there exists an integer $k \geq 0$ such that $s^k m_i$ lies in W for all $1 \leq i \leq r$. In particular, we have $M \subset M_k = \mathrm{Tr}_{H_{\Delta, k}/s^k H_{\Delta, 0} s^{-k}}(s^k M) \subseteq B_+ \{s^k m_1, \dots, s^k m_r\} \subseteq W$ whence we also deduce $M_\infty = B_+ M \subseteq W$. So M_∞ is a minimal generating B_+ -subrepresentation of $\mathrm{gr}_M^{w_0}(\pi)$, hence we have $D_{SV}(\mathrm{gr}_M^{w_0}(\pi)) = M_\infty^\vee$ by definition. \square

Question 2. *In what generality is it true that $D_\Delta^\vee(\mathrm{gr}_M^{w_0}(\pi)) \cong M^\vee[1/X_\Delta]$?*

Remark. The answer to the above question is affirmative if π is a principal series.

4.6 The fully faithful property of D_Δ^\vee for principal series

We denote by SP_A^0 the category of those finite length smooth representations of G over A whose irreducible subquotients are (necessarily irreducible) principal series representations.

Lemma 4.24. *Let χ and χ' be two (not necessarily distinct) characters $T \rightarrow \kappa^\times$ such that both $\mathrm{Ind}_B^G \chi$ and $\mathrm{Ind}_B^G \chi'$ are irreducible. Then the natural map*

$$\mathrm{Ext}_G^1(\mathrm{Ind}_B^G \chi', \mathrm{Ind}_B^G \chi) \rightarrow \mathrm{Ext}^1(D_\Delta^\vee(\mathrm{Ind}_B^G \chi), D_\Delta^\vee(\mathrm{Ind}_B^G \chi'))$$

is injective.

Proof. We distinguish two cases whether $\chi = \chi'$ or not. Assume first $\chi \neq \chi'$. By Thm. 1.1 in [19] if the left hand side is nonzero then we have $\chi' = s_\alpha(\chi) \cdot \varepsilon^{-1} \circ \alpha$ for some simple root $\alpha \in \Delta$ where $s_\alpha(\chi)$ denotes the conjugate of χ with respect to the reflection corresponding to α in the Weyl group $N_G(T)/T$ and ε is the modulo p cyclotomic character. By our assumption that $\mathrm{Ind}_B^G \chi$ is irreducible, we have $s_\alpha(\chi) \neq \chi$ so we can only have $\chi' = s_\alpha(\chi) \cdot \varepsilon^{-1} \circ \alpha$ for at most one element $\alpha \in \Delta$. Moreover, in this case the unique nonsplit extension π comes from parabolic induction from $G_\alpha \cong TGL_2(\mathbb{Q}_p)$ generated by T , N_α , and s_α . Therefore even Breuil's functor $D_\Delta^\vee(\pi) \cong \kappa((X)) \otimes_{\ell, \kappa((N_{\Delta, 0}))} D_\Delta^\vee(\pi)$ (see Cor. 3.20) is a non-split extension of (φ, Γ) -modules by Thm. 6.1 in [8] and Thm. VII.5.2 in [13].

Now assume $\chi = \chi'$. Then by Thm. 1.2 in [19] the natural map

$$\mathrm{Ext}_T^1(\chi, \chi) \rightarrow \mathrm{Ext}_G^1(\mathrm{Ind}_B^G \chi, \mathrm{Ind}_B^G \chi)$$

is bijective. However, the composite map $\mathrm{Ext}_T^1(\chi, \chi) \rightarrow \mathrm{Ext}^1(D_\Delta^\vee(\mathrm{Ind}_B^G \chi), D_\Delta^\vee(\mathrm{Ind}_B^G \chi))$ is injective since whenever δ is an extension of χ by itself then we have the identification $D_\Delta^\vee(\mathrm{Ind}_B^G \delta) \cong \kappa((N_{\Delta, 0})) \otimes \delta^\vee$ and we may recover δ from this by taking the Pontryagin dual of the intersection

$$\bigcap_{n>0} \varphi_s^n(\kappa((N_{\Delta, 0})) \otimes \delta^\vee) \cong \left(\bigcap_{n>0} \varphi_s^n(\kappa((N_{\Delta, 0})) \right) \otimes \delta^\vee \cong \delta^\vee.$$

\square

Remark. We do not know whether or not the above map between extension groups is also surjective. For this one should determine the dimension of the right hand side. The author plans to turn back to this question in a future work.

Lemma 4.25. *Let π be an object in the category SP_A^0 and D be an object in $\mathcal{D}^{et}(T_+, A((N_{\Delta,0})))$ that is a successive extension of rank 1 objects. Assume that the length of π is bigger than that of D . Then π^\vee cannot be embedded into the global sections of a G -equivariant sheaf \mathfrak{Y} with an injection $\mathfrak{Y}(\mathcal{C}_0) \hookrightarrow \mathbb{M}_{\infty,0}(D)$.*

Proof. Assume it can. By Prop. 4.14 and Thm. 4.9 (or simply by Prop. 2.8) the composite map

$$\pi^\vee \hookrightarrow \mathfrak{Y}(G/B) \twoheadrightarrow \mathfrak{Y}(\mathcal{C}_0) \hookrightarrow \mathbb{M}_{\infty,0}(D) \twoheadrightarrow \mathbb{D}_{0,\infty} \circ \mathbb{M}_{\infty,0}(D) \cong D$$

factors uniquely through $D_\Delta^\vee(\pi)$. In particular, there exists an $M \in \mathcal{M}_\Delta^0(\pi^{H_{\Delta,0}})$ such that we have the factorization $\pi^\vee \twoheadrightarrow M^\vee \hookrightarrow M^\vee[1/X_\Delta] \hookrightarrow D$. We may assume without loss of generality that we have $M^\vee[1/X_\Delta] \cong D$. By assumption we have a filtration on π whose subquotients are principal series. By Theorem 3.19 this induces a filtration on $D_\Delta^\vee(\pi)$ whose subquotients are 1-dimensional objects. Further, this induces a filtration on D via the surjective map $D_\Delta^\vee(\pi) \twoheadrightarrow D$ whence we also obtain a filtration on the sheaf \mathfrak{Y} and its global sections $\mathfrak{Y}(G/B)$. Therefore passing to one particular subquotient we may assume without loss of generality that π has length 2 and D is 1 dimensional. So π is an extension of 2 principal series for characters χ and χ' . In particular, we have two surjective maps $D_\Delta^\vee(\pi) \twoheadrightarrow D$ and $D_\Delta^\vee(\pi) \twoheadrightarrow D_\Delta^\vee(\text{Ind}_B^G \chi)$. The kernels of these maps must be different since the map $\pi^\vee \rightarrow \mathfrak{Y}_{\pi, M_\chi}(G/B)$ has $(\text{Ind}_B^G \chi')^\vee$ in its kernel but $\pi^\vee \rightarrow \mathfrak{Y}(G/B)$ is injective. (Here M_χ denotes the unique element in $\mathcal{M}_\Delta^0((\text{Ind}_B^G \chi)^{H_{\Delta,0}})$.) In particular, $D_\Delta^\vee(\pi)$ splits as a direct sum of these two kernels whence by Lemma 4.24 $\pi \cong \text{Ind}_B^G \chi \oplus \text{Ind}_B^G \chi'$. Therefore the projection to D has kernel $D_\Delta^\vee(\text{Ind}_B^G \chi)$ that contradicts to the assumption that $\pi^\vee \rightarrow \mathfrak{Y}(G/B)$ is injective. \square

Our main result in this section is the following

Theorem 4.26. *The restriction of D_Δ^\vee to the category SP_A^0 is fully faithful.*

Proof. The faithfulness is clear by the exactness (Thm. 3.19) noting that D_Δ^\vee does not vanish on irreducible objects in SP_A^0 by Cor. 3.9. Let π and π' be objects in the category SP_A^0 and assume we have a nonzero morphism $f: D_\Delta^\vee(\pi) \rightarrow D_\Delta^\vee(\pi')$. Let D be the image of f and n be the (generic) length of D . Consider the sheaf $\mathfrak{Y}_{\pi, \Delta}$ on G/B provided by Cor. 4.22. By Prop. 3.18 $\mathfrak{Y}_{\pi, \Delta} = \mathfrak{Y}_{\pi, M}$ for some M in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$ and we have $\text{length}_{gen}(M^\vee[1/X_\Delta]) = \text{length}(\pi)$. Further, the image of the composite map $M^\vee \rightarrow M^\vee[1/X_\Delta] = D_\Delta^\vee(\pi) \xrightarrow{f} D$ is the Pontryagin dual of another object $M_1 \leq M$ in $\mathcal{M}_\Delta(\pi^{H_{\Delta,0}})$. Since f is onto, $D = M_1^\vee[1/X_\Delta]$. By 4.21 we obtain a morphism $\mathfrak{Y}_{\pi, M} \rightarrow \mathfrak{Y}_{\pi, M_1}$ of G -equivariant sheaves whose kernel has sections

$$\mathfrak{Y}_{\pi, \text{Ker}(f)}(U) := \{x \in \mathfrak{Y}_{\pi, M}(U) \mid \text{res}_{gU \cap \mathcal{C}_0}^{gU}(gx) \in \text{Ker}(f) \subseteq M^\vee[1/X_\Delta] \text{ for all } g \in G\}$$

on an open subset $U \subseteq G/B$. Let $f_{G/B}: \pi^\vee \rightarrow \mathfrak{Y}_{\pi, M_1}(G/B)$ be the natural map and π_1^\vee be its image and π_2^\vee be its kernel. Then $\pi_1 := (\pi_1^\vee)^\vee$ is naturally a G -subrepresentation of π with quotient $\pi_2 = (\pi_2^\vee)^\vee$. By Lemma 4.25 applied to both $\pi_1^\vee \hookrightarrow \mathfrak{Y}_{\pi, M_1}(G/B)$ and $\pi_2^\vee \hookrightarrow \mathfrak{Y}_{\pi, \text{Ker}(f)}(G/B)$ we obtain $\text{length}(\pi_1) \leq n$ and $\text{length}(\pi_2) \leq \text{length}_{gen}(\text{Ker}(f)) = \text{length}(\pi) - n$. Hence $\text{length}(\pi_1) = n$ and $D_\Delta^\vee(\pi_1) \cong D$.

By a similar argument applied to the quotient map $D_\Delta^\vee(\pi') \twoheadrightarrow \text{Coker}(f)$ we find a quotient $\pi' \twoheadrightarrow \pi'_2$ with an injective morphism $\pi_2'^\vee \hookrightarrow \mathfrak{Y}_{\pi, D}(G/B)$ into the global sections of a sheaf $\mathfrak{Y}_{\pi, D}$ whose sections on an open subset $U \subseteq G/B$ are given by

$$\mathfrak{Y}_{\pi, D}(U) := \{x \in \mathfrak{Y}_{\pi', \Delta}(U) \mid \text{res}_{gU \cap \mathcal{C}_0}^{gU}(gx) \in D \subseteq D_\Delta^\vee(\pi') \text{ for all } g \in G\}.$$

Further, we have $\text{length}(\pi'_2) = n$ and $D_\Delta^\vee(\pi'_2) \cong D$.

Now we have maps $\pi_1^\vee \rightarrow D_\Delta^\vee(\pi_1) \cong D$ and $\pi_2^{\vee'} \rightarrow D_\Delta^\vee(\pi_2) \cong D$ so we may take the direct sum $\theta: \pi_1^\vee \oplus \pi_2^{\vee'} \rightarrow D$. This gives rise to an object $M_* \in \mathcal{M}_\Delta((\pi_1 \oplus \pi_2)^{H_{\Delta,0}})$ and a G -equivariant continuous map $\bar{\theta}: \pi_1^\vee \oplus \pi_2^{\vee'} \rightarrow \mathfrak{Y}_{\pi_1 \oplus \pi_2, M_*}(G/B)$ that is injective restricted to both π_1^\vee and to $\pi_2^{\vee'}$. However, applying Lemma 4.25 to this situation again we deduce that the image of $\bar{\theta}$ has length at most n . Therefore $\bar{\theta}$ induces an isomorphism between π_1^\vee and $\pi_2^{\vee'}$ (and the image). All in all we obtain a map $\pi' \rightarrow \pi'_1 \cong \pi_1 \hookrightarrow \pi$ that induces f after taking D_Δ^\vee . \square

Remarks. 1. In particular, the forgetful functor restricting π to B is also fully faithful on SP_A^0 as D_Δ^\vee factors through this.

2. D_Δ^\vee is not faithful on SP_A : for instance any finite dimensional representation lies in its kernel.

References

- [1] Berger L., Représentations modulaires de $\text{GL}_2(\mathbb{Q}_p)$ et représentations galoisiennes de dimension 2, *Astérisque* **330** (2010), 263–279.
- [2] Berger L., Multivariable Lubin-Tate (φ, Γ) -modules and filtered φ -modules, *Math. Res. Lett.* **20** (2013), no. 3, 409–428.
- [3] Berger L., Multivariable (φ, Γ) -modules and locally analytic vectors, preprint (2016), to appear in *Duke Math. Journal*
- [4] Breuil Ch., Sur quelques représentations modulaires et p -adiques de $\text{GL}_2(\mathbb{Q}_p)$ I, *Compositio Math.* **138** (2003), 165–188.
- [5] Breuil Ch., Sur quelques représentations modulaires et p -adiques de $\text{GL}_2(\mathbb{Q}_p)$ II, *J. Inst. Math. Jussieu* **2** (2003), 1–36.
- [6] Breuil Ch., Emerton M., Représentations p -adiques ordinaires de $\text{GL}_2(\mathbb{Q}_p)$ et compatibilité local-global, *Astérisque* **331** (2010), 255–315.
- [7] Breuil Ch., The emerging p -adic Langlands programme, in: *Proceedings of the International Congress of Mathematicians Volume II*, Hindustan Book Agency, New Delhi (2010), 203–230.
- [8] Breuil Ch., Induction parabolique et (φ, Γ) -modules, *Algebra & Number Theory* **9**(10) (2015), 2241–2291.
- [9] Breuil Ch., private communication, 11/03/2016
- [10] Breuil Ch., Herzig F., Ordinary representations of $G(\mathbb{Q}_p)$ and fundamental algebraic representations, *Duke Math. Journal* **164** (2015), 1271–1352.
- [11] Breuil Ch., V. Paškūnas, Towards a modulo p Langlands correspondence for GL_2 , *Memoirs of the AMS*, **216** (2012).

- [12] Colmez P., (φ, Γ) -modules et représentations du mirabolique de $GL_2(\mathbb{Q}_p)$, *Astérisque* **330** (2010), 61–153.
- [13] Colmez P., Représentations de $GL_2(\mathbb{Q}_p)$ et (φ, Γ) -modules, *Astérisque* **330** (2010), 281–509.
- [14] Emerton M., On a class of coherent rings, with applications to the smooth representation theory of $GL_2(\mathbb{Q}_p)$ in characteristic p , preprint (2008).
- [15] Emerton M., Ordinary parts of admissible representations of p -adic reductive groups I. Definition and first properties, *Astérisque* **331** (2010), 355–402.
- [16] Erdélyi M., On the Schneider–Vigneras functor for principal series, *J. of Number Theory* **162** (2016), 68–85.
- [17] Erdélyi M., Zábrádi G., Links between generalized Montréal functors, preprint (2015), arXiv:1412.5778
- [18] Fontaine J.-M., Représentations p -adiques des corps locaux, in “The Grothendieck Festschrift”, vol 2, Prog. in Math. 87, 249–309, Birkhäuser 1991.
- [19] Hauseux J., Complements sur les extensions entre series principales p -adiques et modulo p de $G(F)$, preprint (2014), arXiv:1407.4630
- [20] Kedlaya K., Some slope theory for multivariate Robba rings, preprint (2013), arXiv:1311.7468
- [21] Kisin M., Deformations of $G_{\mathbb{Q}_p}$ and $GL_2(\mathbb{Q}_p)$ representations, *Astérisque* **330** (2010), 511–528.
- [22] Li H., van Oystaeyen F., *Zariskian Filtrations*, *K*-monographs in Mathematics, vol. 2, Kluwer Academic Publishers, Dordrecht, 1996.
- [23] McConnell J. C., Robson J. C., *Noncommutative Noetherian Rings*, Graduate Studies in Mathematics, vol. **30**, AMS, Providence, Rhode Island, 1987.
- [24] Paškūnas V., The image of Colmez’s Montreal functor, *Publ. Math. IHES* **118**(1) (2013), 1–191.
- [25] Schneider P., Vigneras M.-F., A functor from smooth \mathfrak{o} -torsion representations to (φ, Γ) -modules, Volume in honour of F. Shahidi, Clay Mathematics Proceedings Volume **13** (2011), 525–601.
- [26] Schneider P., Vigneras M.-F., Zábrádi G., From étale P_+ -representations to G -equivariant sheaves on G/P , in: *Automorphic forms and Galois representations* (Volume 2), LMS Lecture Note Series **415** (2014), Cambridge Univ. Press, 248–366.
- [27] Vigneras M.-F., Série principale modulo p de groupes réductifs p -adiques, *Geom. and Funct. Analysis* **17** (2008), 2090–2112.

- [28] Zábrádi G., (φ, Γ) -modules over noncommutative overconvergent and Robba rings, *Algebra & Number Theory* **8**(1) (2014), 191–242.
- [29] Zábrádi G., Multivariable (φ, Γ) -modules and products of Galois groups, preprint (2016)